

Haploid (2,2)-Superfields In 2-Dimensional Spacetime

Tristan Hübsch♣

*Department of Physics and Astronomy
Howard University, Washington, DC 20059
thubsch@howard.edu*

ABSTRACT

Superfields in 2-dimensional (2,2)-superspacetime which are independent of (some) half of the fermionic coordinates are discussed in a hopefully both comprehensive and comprehensible manner. An embarrassing abundance of these simplest ‘building blocks’ makes it utterly impossible to write down the ‘most general Lagrangian’. With some *ad hoc* but perhaps plausible restrictions, a rather general Lagrangian is found, which exhibits many of the phenomena that have been studied recently, and harbors many more. In particular, it becomes patently obvious that the (2,2)-supersymmetric 2-dimensional field theory target space geometries (many of which are suitable for (super)string propagation) are far more general than Kähler manifolds with holomorphic bundles.

December '98.

♣ On leave from the “Rudjer Bošković” Institute, Zagreb, Croatia.

1. Introduction

Superfields are used extensively in 2-dimensional supersymmetric field theories. Alas! the diversity of conventions obscures the straightforward. In particular, these building blocks are frequently obtained by dimensional reduction from the 4-dimensional case, which (by far) does not provide for full generality [1]: contrary to some claims in the literature (even in book form!) in 2-dimensional spacetimes, *(2,2)- (a.k.a. $N=2$) supersymmetry does not imply Kählerness of the field space!*

An intrinsically 2-dimensional analysis, ‘from scratch’, is presented here, and provides generalizations to the results in the recent literature; see, *e.g.*, Refs. [2,3,4,5].

This article is organized as follows: The remaining part of this introductory section presents some basic facts about 2-dimensional (2,2)-superspace and sets up the notation; further details are found in Appendix A. Section 2 presents a rather rich set of constrained superfields in a systematic way, and provides several directions for generalizations. The ‘most general’ Lagrangian for the superfields of §2 (subject to a few hopefully plausible restrictions) is given in Section 3, using ‘implicitly constrained’ superfields, with the details of component field expansions deferred to the Appendix B. An alternate, ‘explicitly constrained’ formalism is discussed in Section 4. Section 5 then turns to some relations to geometry, in particular that of the target (field) space. Section 6 summarizes the presented results and discusses some further topics.

1.1. 2-dimensional spacetime

The crucial peculiarity of supersymmetric models in 2-dimensional spacetime stems from the fact that the Lorentz group, $SO(1,1)$, is Abelian and so has only 1-dimensional irreducible representations: for example, the coordinate 2-vector (σ^0, σ^1) decomposes into the (light-cone) characteristic coordinates $\sigma^{\pm\pm} \stackrel{\text{def}}{=} \frac{1}{2}(\sigma^0 \pm \sigma^1)$. These, in fact, are eigenfunctions of the (Lorentz boost) generator¹⁾, $\mathbf{B} \stackrel{\text{def}}{=} (\sigma^1 \partial_0 + \sigma^0 \partial_1)$, of $SO(1,1)$:

$$\mathbf{B} \sigma^{\pm\pm} = j_3(\sigma^{\pm\pm}) \sigma^{\pm\pm} = \pm \sigma^{\pm\pm} \quad (1.1)$$

so that the group element, $\mathbf{U}_\alpha \stackrel{\text{def}}{=} e^{i\alpha \mathbf{B}}$, acts diagonally: $\mathbf{U}_\alpha \sigma^{\pm\pm} = e^{\pm i\alpha} \sigma^{\pm\pm}$. For brevity, we also write $\sigma^{\pm} \stackrel{\text{def}}{=} \sigma^{++}$ and $\sigma^= \stackrel{\text{def}}{=} \sigma^{--}$. Note that upon the analytic continuation $\sigma^0 \rightarrow i\sigma^0$, $(\sigma^{\pm}, \sigma^=) \rightarrow (z, -\bar{z})$: the light-cone structure becomes a complex structure. Similar arguments show that all other tensors (spinors) decompose into 1-component objects. Upon the analytic continuation to imaginary time, $\sigma^0 \rightarrow i\sigma^0$, $\mathbf{U}_\alpha$ becomes the winding number (holomorphic homogeneity) operator. The sub- and superscripts “ $\pm$ ” then simply denote the winding number (spin in real time) in units of $\frac{1}{2}(\hbar)$. Functions depending on only $\sigma^{\pm}$ or only $\sigma^=$ are called left- and right-movers, respectively, and become holomorphic (complex-analytic) and anti-holomorphic (complex-antianalytic) functions upon analytic continuation to imaginary time.

¹⁾ The eigenvalue of the Lorentz boost operator, $\mathbf{B} \stackrel{\text{def}}{=} (\sigma^1 \partial_0 + \sigma^0 \partial_1)$, extended to total angular momentum in the usual way, equals the spin projection, and will therefore be denoted by j_3 although it does not stem from its 4-dimensional namesake. Moreover, since all representations of the 2-dimensional Lorentz group are 1-dimensional, there is no need to distinguish between ‘spin’ and ‘spin projection’, whence we will call j_3 simply ‘spin’.

1.2. (2,2)-superbasics

The (2,2)-supersymmetry algebra involves the supersymmetry charges $Q_{\pm}$ and $\bar{Q}_{\pm}$, which satisfy (adapting from Refs. [6,2]; comparison with Refs. [7,4] is provided in appendix A):

$$\{Q_{\pm}, \bar{Q}_{\pm}\} = -2i\partial_{\pm\pm} = 2(H \pm p). \quad (1.2)$$

Here $\partial_{++} \equiv \partial_{+} \stackrel{\text{def}}{=} \frac{\partial}{\partial \sigma^{+}} = (\partial_0 + \partial_1)$ and $\partial_{--} \equiv \partial_{-} \stackrel{\text{def}}{=} \frac{\partial}{\partial \sigma^{-}} = (\partial_0 - \partial_1)$. $p = -i\partial_1 = -\frac{i}{2}(\partial_{+} - \partial_{-})$ is the linear momentum operator and $H = -i\partial_0 = -\frac{i}{2}(\partial_{+} + \partial_{-})$ is the Hamiltonian (energy) operator. Note that p has the usual sign as in Quantum Mechanics, while H has the opposite sign; with these conventions, $(-E, E)$ is a light-like 2-vector.

Equipping the world-sheet with anticommuting fermionic coordinates, $\varsigma^{\pm}, \bar{\varsigma}^{\pm}$, the supercharges are realized as differential operators on the (super) world-sheet²⁾:

$$Q_{\pm} \stackrel{\text{def}}{=} \partial_{\pm} + i\bar{\varsigma}^{\pm}\partial_{\pm\pm}, \quad \bar{Q}_{\pm} \stackrel{\text{def}}{=} -\bar{\partial}_{\pm} - i\varsigma^{\pm}\partial_{\pm\pm}. \quad (1.3)$$

Supersymmetry transformations are generated by the $Q, \bar{Q}$'s, and implemented (effected) by unitary operator

$$U_{\epsilon, \bar{\epsilon}} \stackrel{\text{def}}{=} \exp \{i(\epsilon^{\pm}Q_{\pm} + \bar{\epsilon}^{\pm}\bar{Q}_{\pm})\}. \quad (1.4)$$

which, in fact, is a quartic multinomial in the $Q, \bar{Q}$'s, by virtue of their nilpotency. The spinorial derivatives

$$D_{\pm} \stackrel{\text{def}}{=} \partial_{\pm} - i\bar{\varsigma}^{\pm}\partial_{\pm\pm}, \quad \bar{D}_{\pm} \stackrel{\text{def}}{=} -\bar{\partial}_{\pm} + i\varsigma^{\pm}\partial_{\pm\pm}, \quad (1.5)$$

are covariant with respect to $U_{\epsilon, \bar{\epsilon}}$. Easily, $\{Q_{\pm}, D_{\pm}\} = 0 = \{\bar{Q}_{\pm}, D_{\pm}\}$, whereupon $[U_{\epsilon, \bar{\epsilon}}, D_{\pm}] = 0 = [U_{\epsilon, \bar{\epsilon}}, \bar{D}_{\pm}]$.

Finally the $D, \bar{D}$'s close virtually the same algebra (1.2), as do the $Q, \bar{Q}$'s:

$$\{D_{\pm}, \bar{D}_{\pm}\} = 2i\partial_{\pm\pm} = -2(H \pm p). \quad (1.6)$$

All anticommutators among the $Q_{\pm}, \bar{Q}_{\pm}, D_{\pm}, \bar{D}_{\pm}$, other than (1.2) and (1.6), vanish.

The Berezin superintegrals (over supercoordinates) are by definition equivalent to partial superderivatives, and up to total (world-sheet) spacetime derivatives³⁾ (which we ignore, assuming world-sheets without boundaries) equivalent to covariant superderivatives [6,7]. We use:

$$\int d^4\varsigma [\dots] \stackrel{\text{def}}{=} \frac{1}{4}[D_{-}, \bar{D}_{-}][D_{+}, \bar{D}_{+}][\dots] \equiv D^4[\dots], \quad (1.7)$$

where “|” means setting $\varsigma^{\pm} = 0 = \bar{\varsigma}^{\pm}$. Integration over a fermionic subspace is formally achieved by inserting a fermionic Dirac delta-function, so that:

$$\int d^2\varsigma [\dots] \equiv \frac{1}{2}[D_{-}, D_{+}][\dots], \quad \int d^2\bar{\varsigma} [\dots] \equiv \frac{1}{2}[\bar{D}_{+}, \bar{D}_{-}][\dots]; \quad (1.8a, b)$$

²⁾ Since $(\partial_{\pm\pm})^{\dagger} = -\partial_{\pm\pm}$, for $(Q_{\pm})^{\dagger} = \bar{Q}_{\pm}$, it must be that $(\varsigma^{\pm})^{\dagger} = -\bar{\varsigma}^{\pm}$ and $(\partial_{\pm})^{\dagger} = -\bar{\partial}_{\pm}$.

³⁾ Hereafter, ‘total derivative’ will stand for ‘total (world-sheet) spacetime derivative’.

$$\int d^2\zeta [\dots] \equiv \frac{1}{2}[D_-, \bar{D}_+][\dots] , \quad \int d^2\bar{\zeta} [\dots] \equiv \frac{1}{2}[D_+, \bar{D}_-][\dots] ; \quad (1.8c, d)$$

$$\int d^2\zeta^+ [\dots] \equiv \frac{1}{2}[D_-, \bar{D}_-][\dots] , \quad \int d^2\zeta^- [\dots] \equiv \frac{1}{2}[D_+, \bar{D}_+][\dots] . \quad (1.8e, f)$$

Clearly, the definitions (1.8c, d) may also be obtained from (1.8a, b) by swapping $\zeta^+ \leftrightarrow \bar{\zeta}^+$; see below. Similarly, the definitions (1.8e, f) follow from (1.8c, d) upon $\bar{\zeta}^+ \leftrightarrow \bar{\zeta}^-$. As expected then, for example,

$$\int d^4\zeta [\dots] = \int d^2\zeta \int d^2\bar{\zeta} [\dots] , = - \int d^2\zeta \int d^2\bar{\zeta} [\dots] . \quad (1.9)$$

Note the negative sign in the second identity, and that the integral measures in (1.8e, f) are not (pseudo)scalars; they have spin ± 1 . Also, a few *operatorial* identities collected in Appendix A will be useful.

1.3. Superspace symmetries

In more than 2-dimensional spacetimes, Lorentz transformations leave the integrals (1.8) unchanged. However, they rotate Eqs. (1.8c) $\leftrightarrow$ (1.8d) and (1.8e) $\leftrightarrow$ (1.8f).

It is of interest then to examine the discrete symmetries acting on the four-fermionic coordinate system, or alternatively, the four covariant superdifferentials, $D_-, \bar{D}_-, D_+, \bar{D}_+$. Since the Lorentz transformations act diagonally on the $D, \bar{D}$'s, they can be treated as independent objects in a Lorentz-invariant fashion. Thus, the maximal discrete group acting on the $D_\pm, \bar{D}_\pm$ system by swapping them is S_4 , the group of permutations of four distinguishable objects. The simplest are, of course, the swaps:

$$C_+ : D_+ \leftrightarrow \bar{D}_+ , \quad C_- : D_- \leftrightarrow \bar{D}_- , \quad (1.10a, b)$$

$$p : D_+ \leftrightarrow D_- , \quad \bar{p} : \bar{D}_+ \leftrightarrow \bar{D}_- , \quad (1.10c, d)$$

$$q : \bar{D}_- \leftrightarrow D_+ , \quad \bar{q} : D_- \leftrightarrow \bar{D}_+ . \quad (1.10e, f)$$

where each of these operations only swaps the indicated superderivatives and leaves the other two intact. Also, it should be clear that $[C_+, C_-]=0$, $[p, \bar{p}]=0$ and $[q, \bar{q}]=0$. Furthermore, *e.g.*, $p C_- p = C_- p C_- = \bar{q}$, and $\bar{p} C_+ \bar{p} = C_+ \bar{p} C_+ = \bar{q}$.

There are additional discrete transformations involving time-reversal, $\mathbf{T}$, which acts by $\mathbf{T}(\partial_+) = -\partial_-$ and $\mathbf{T}(\partial_-) = -\partial_+$. Since time-reversal squares to the identity, $\mathbf{T}^2 = \mathbb{1}$, it will 'double' the group of discrete transformations of the $D_\pm, \bar{D}_\pm$ system, and we call this group $2S_4$.

Notice that the familiar (ambidexterous) complex conjugation, $\mathbf{C}$, may be expressed as $\mathbf{C} = C_- C_+$. This operation exchanges the integrals (1.8a) $\leftrightarrow$ (1.8b) and (1.8c) $\leftrightarrow$ (1.8d), but leaves (1.8e) and (1.8f) intact.

Similarly, the full parity (spatial reflection), $\mathbf{P}$, may be expressed as $\mathbf{P} = p\bar{p}$, and it is easy to show that $[\mathbf{C}, \mathbf{P}]=0$, as should be the case. $\mathbf{P}$ exchanges the integrals (1.8c) $\leftrightarrow$ (1.8d) and (1.8e) $\leftrightarrow$ (1.8f), but leaves (1.8a) and (1.8b) intact.

Also, the combined parity and complex conjugation, $\mathbf{CP} = \mathbf{q}\bar{\mathbf{q}}$, exchanges the integrals (1.8a) $\leftrightarrow$ (1.8b) and (1.8e) $\leftrightarrow$ (1.8f), but leaves those in (1.8c) and (1.8d) intact.

It is very important to realize that the operators (1.10) are defined there to act on the ‘soul’ of the (2,2)-super-Riemann surface, *i.e.*, the fermionic coordinates and so the superderivatives acting on them. However, the relation between the superderivatives and the world-sheet derivatives (1.6) induces an action on the ‘body’ too, *i.e.*, on the world-sheet coordinates. In the often used Euclidean incarnation, $(\sigma^\pm, \sigma^\pm) \rightarrow (z, -\bar{z})$, what was called ‘parity’ (the unidexterous $\mathbf{p}$ and $\bar{\mathbf{p}}$, and the ambidexterous $\mathbf{P}$) becomes (another!) complex conjugation. While the induced action of $\mathbf{C}_\pm$ and $\mathbf{C}$ on the world-sheet coordinates $(\sigma^\pm, \sigma^\pm)$ is trivial (since the supersymmetry relations (1.6) are Hermitian), the induced action of $\mathbf{p}$, $\bar{\mathbf{p}}$ and $\mathbf{P}$ is not! These do become the (unidexterous and ambidexterous, respectively) complex conjugations when acting on the Euclideanized ‘body’, *i.e.*, bosonic world-sheet coordinates, but remain parity (spin-flip) in the ‘soul’, *i.e.*, the fermionic coordinates. Foremost, they are independent from the $\mathbf{C}_\pm$ and $\mathbf{C}$. The unaware Reader should be cautioned that the careful distinction between these operations is sometimes muddled in the literature.

Finally, the Lorentz boost group element (generalized, as usual, to include spin) $\mathbf{U}_\alpha = e^{i\alpha\mathbf{B}}$ satisfies $\mathbf{U}_\alpha\mathbf{U}_\beta = \mathbf{U}_{(\alpha+\beta)}$ so $[\mathbf{U}_\alpha, \mathbf{U}_\beta] = 0$. Also, $[\mathbf{U}_\alpha, \mathbf{C}_\pm] = 0$, but $\mathbf{U}_\alpha\mathbf{P} = \mathbf{P}\mathbf{U}_\alpha^{-1}$. Thus, the usual super-Poincaré group is extended by the various discrete symmetries (1.10) and time-reversal, $\mathbf{T}$, so that the discrete subgroup is $2S_4$.

The discrete symmetry, S_4 , generated by the swaps (1.10), is part of the continuous $GL(4, \mathbb{C})$, the $GL_-(2, \mathbb{C}) \times GL_+(2, \mathbb{C})$ subgroup of which then commutes with the Lorentz symmetry. I defer the study of such continuous transformations to a later opportunity.

2. Haploid Superfields

General superfields are (super)functions of $(\sigma^0, \sigma^1; \varsigma^\pm, \bar{\varsigma}^\pm)$. Akin to complex-analytic (holomorphic) functions depending only on the complex-analytic half of complex variables, we define *haploid* superfields to depend only on (some) half of the four fermionic coordinates $\varsigma^\pm, \bar{\varsigma}^\pm$. Models that involve such ‘super-holomorphic’ superfields exhibit a much better quantum behavior, not unrelated to the special features of complex-analytic functions [6,7].

2.1. First-order constrained haploid superfields

As the Lorentz transformations in 1+1-dimensional spacetime leave the four $\varsigma^\pm, \bar{\varsigma}^\pm$ independent of each other, there exist *six* distinct types of hapliod superfields (as $\binom{4}{2} = 6$), all of which are annihilated by (some) half of the superderivatives:

$$1. \text{ Chiral : } \quad \bar{D}_+\Phi = 0 = \bar{D}_-\Phi , \quad (2.1a)$$

$$2. \text{ Antichiral : } \quad D_+\bar{\Phi} = 0 = D_-\bar{\Phi} , \quad (2.1b)$$

$$3. \text{ Twisted-chiral : } \quad D_+\Xi = 0 = \bar{D}_-\Xi , \quad (2.1c)$$

$$4. \text{ Twisted-antichiral : } \quad \bar{D}_+\bar{\Xi} = 0 = D_-\bar{\Xi} , \quad (2.1d)$$

$$5. \text{ Lefton : } \quad D_-\Lambda = 0 = \bar{D}_-\Lambda , \quad (2.1e)$$

$$6. \text{ Righton : } \quad D_+\Upsilon = 0 = \bar{D}_+\Upsilon . \quad (2.1f)$$

The first four pairs of superconstraints simply restrict $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}$ to be independent of two of the fermionic coordinates $\varsigma, \bar{\varsigma}$, up to total derivatives.

By contrast, of the last two superconstraint pairs, each one *includes* also a spacetime derivative constraint (see below):

$$\text{Eqs. (2.1e)} \ni \{ \partial_{=} \Lambda = 0 \} , \quad \text{Eqs. (2.1f)} \ni \{ \partial_{\pm} \Upsilon = 0 \} . \quad (2.2)$$

As a result, the component fields in the lepton superfield Λ are functions of only $\sigma^{\pm}$ ('holomorphic', *i.e.*, 'left-movers'), whereas those in the righton superfield Υ are functions of only $\sigma^{\pm}$ ('anti-holomorphic', *i.e.*, 'right-movers'). We return to this below, after defining the component fields.

We will also encounter an even more restricted type of superfields, which depend only on one out of the four $\varsigma, \bar{\varsigma}$'s, and so are annihilated by *three* superderivatives:

$$1. \text{ Chiral Lefton : } \quad \bar{D}_{+} \Phi_L = \bar{D}_{-} \Phi_L = D_{-} \Phi_L = 0 , \quad (2.3a)$$

$$2. \text{ Antichiral Lefton : } \quad D_{+} \bar{\Phi}_L = \bar{D}_{-} \bar{\Phi}_L = D_{-} \bar{\Phi}_L = 0 , \quad (2.3b)$$

$$3. \text{ Chiral Righton : } \quad \bar{D}_{-} \Phi_R = \bar{D}_{+} \Phi_R = D_{+} \Phi_R = 0 , \quad (2.3c)$$

$$4. \text{ Antichiral Righton : } \quad D_{-} \bar{\Phi}_R = \bar{D}_{+} \bar{\Phi}_R = D_{+} \bar{\Phi}_R = 0 . \quad (2.3d)$$

Depending only on some quarter of the supercoordinates $\varsigma, \bar{\varsigma}$, we call them *quartoid* superfields. Clearly, they may be considered as special cases of the haploid ones (2.1): *e.g.*, Φ_L may be regarded as a D_{-} -annihilated chiral superfield, a $\bar{D}_{-}$ -annihilated twisted antichiral superfield, or a $\bar{D}_{+}$ -annihilated (*i.e.*, chiral) lepton. We adopted this last nomenclature in virtue of its simplicity and the fact that it also alludes to the implicit spacetime constraints $\partial_{=} \Phi_L = 0 = \partial_{=} \bar{\Phi}_L$ and $\partial_{\pm} \Phi_R = 0 = \partial_{\pm} \bar{\Phi}_R$. The Reader may prefer the obvious twisted analogues: $\Xi_L = \bar{\Phi}_L$, $\Xi_R = \Phi_R$, $\bar{\Xi}_L = \Phi_L$ and $\bar{\Xi}_R = \bar{\Phi}_R$.

A disclaimer is in order: our purpose thus far was merely to list the logical possibilities. The quantum theory of the unidexterous fields defined in Eqs. (2.1e, f) and (2.3a–d) may indeed seem curiously cumbersome, owing to the occurrence of the spacetime constraints as exemplified in Eqs. (2.2). For unidexterous spin-0 fields ('chiral bosons') and their (1,0)- and (1,1)-supersymmetrizations see Refs. [8,9]; Ref. [10] offers some welcome clarifications about the dynamics and symmetries involved and an updated reference list on the topic. We merely remark that the constraints may be regarded as generators of a gauge symmetry, with the transformations:

$$\begin{aligned} \partial_{=} \Lambda = 0 & \rightarrow \delta \Lambda = \kappa^{\pm} (\partial_{=} \Lambda) , \\ \partial_{\pm} \Upsilon = 0 & \rightarrow \delta \Upsilon = \kappa^{\pm} (\partial_{\pm} \Upsilon) . \end{aligned} \quad (2.4)$$

This gauge invariance then ensures the proper decoupling of the unwanted degrees of freedom. Thus, a well-defined quantum field theory model involving unidexterous (super)fields must be invariant with respect to the chiral gauge symmetry (2.4).

Seeking generality, our approach will then be as follows: at first, we ignore the unidexterous quantization issues. We then return to these issues in a simple setting and discuss some of the implications. Relying on the general and powerful machineries such as the operatorial quantization (see Ref. [11] and the references therein), we defer the (careful tedium of *Ausarbeitung* of the) covariantization of (a suitable subset of) the present results with respect to (2.4) and a full treatment of the (super)constraint effects to a later effort.

2.2. Superfield symmetries

In more than 2-dimensional spacetimes, Lorentz transformations leave the pair of superconstraint superderivatives $(2.1a, b)$ (and so also $\Phi, \bar{\Phi}$) unchanged. However, they mix those in Eqs. $(2.1c) \leftrightarrow (2.1d)$ and $(2.1e) \leftrightarrow (2.1f)$. So, for example, requiring a superfield to satisfy Eq. $(2.1c)$ implies, upon a suitable Lorentz transformation, that *the same* superfield must also satisfy $(2.1d)$, whereupon it follows that it must be a constant. The same applies to the Eqs. $(2.1e, f)$. Only chiral superfields and their conjugates remain viable haploid superfields when spacetime has more than two dimensions. Conversely, $\Xi, \bar{\Xi}, \Lambda, \Upsilon$ cannot be obtained by dimensional reduction; their existence is exceptional to $(2,2)$ -supersymmetric $(\leq)2$ -dimensional spacetime.

Now, complex conjugation⁴⁾, $\mathbf{C}$, leaves the superconstraints $(2.1e, f)$ invariant, while exchanging $(2.1a) \leftrightarrow (2.1b)$ and $(2.1c) \leftrightarrow (2.1d)$, and so $\Phi \leftrightarrow \bar{\Phi}$ and $\Xi \leftrightarrow \bar{\Xi}$. Thus, $\Phi, \bar{\Phi}$ and $\Xi, \bar{\Xi}$ are complex conjugate pairs, while Λ and Υ remain unrelated. So, while in general complex, the unidexterous superfields Λ and Υ *may be chosen* to be real or imaginary. Note that the ‘left-handed conjugation’, $\mathbf{C}_+$, exchanges $\Phi \leftrightarrow \Xi$ and $\bar{\Phi} \leftrightarrow \bar{\Xi}$, while the ‘right-handed conjugation’, $\mathbf{C}_-$ exchanges $\Phi \leftrightarrow \bar{\Xi}$ and $\bar{\Phi} \leftrightarrow \Xi$, and both leave Λ, Υ intact.

On the other hand, parity, $\mathbf{P}$, reflects spin $(+ \leftrightarrow -)$. Therefore, it exchanges Eqs. $(2.1c) \leftrightarrow (2.1d)$ and $(2.1e) \leftrightarrow (2.1f)$, and so $\Xi \leftrightarrow \bar{\Xi}$ and $\Lambda \leftrightarrow \Upsilon$, but leaves Eqs. $(2.1a, b)$ unchanged. That is, $\Xi, \bar{\Xi}$ and Λ, Υ are paired by the left-right parity, $\mathbf{P}$, while Φ and $\bar{\Phi}$ remain unrelated. Note that the ‘unconjugate parity’, $\mathbf{p}$, exchanges $\Xi \leftrightarrow \Lambda$ and $\bar{\Xi} \leftrightarrow \Upsilon$, while the ‘conjugate parity’, $\bar{\mathbf{p}}$ exchanges $\Xi \leftrightarrow \Upsilon$ and $\bar{\Xi} \leftrightarrow \Lambda$, and both leave $\Phi, \bar{\Phi}$ intact.

However, since $\mathbf{C}$ commutes with $\mathbf{P}$, in $\mathbf{C}$ -invariant models Φ and $\bar{\Phi}$ must transform identically under the action of $\mathbf{P}$ and any other symmetry that commutes with $\mathbf{C}$. Similarly, in ‘left-right-symmetric’ (parity-invariant) models, Λ and Υ must transform identically under the action of $\mathbf{C}$ and any other symmetry that commutes with $\mathbf{P}$.

Besides the S_4 group of discrete symmetries (1.10) and time-reversal $\mathbf{T}$, there may also exist discrete or continuous transformations operating entirely in the (super)field space. The fact that the unidexterous superfields $(2.1e, f)$ and (2.3) obey a world-sheet spacetime constraint sets them apart from the others $(2.1a-d)$, and there can be no transformation in the superfield space which would mix the ambidexterous superfields $(2.1a-d)$ and the unidexterous ones $(2.1e, f)$ and (2.3) . This also means that we cannot require of models to be $\mathbf{p}$,- $\bar{\mathbf{p}}$,- $\mathbf{q}$,- or $\bar{\mathbf{q}}$ -invariant. Further symmetry considerations (and especially gauging) are beyond the scope of this paper, but we note that $\mathbf{C}$ -invariance ensures that $(\Phi, \bar{\Phi})$, $(\Xi, \bar{\Xi})$, $(\Phi_L, \bar{\Phi}_L)$ and $(\Phi_R, \bar{\Phi}_R)$ all come in (conjugate) pairs. It is $\mathbf{C}_+$ - (or $\mathbf{C}_-$ -) invariance that would also require that the (anti)chiral and twisted (anti)chiral superfields are paired. No such restriction ensues for the unidexterous superfields Λ, Υ .

⁴⁾ In 2-dimensional spacetime, hermitian conjugation is complex conjugation and transposition (including derivative action to the left), *without* parity or time-reversal.

2.3. Component fields

The fermionic coordinates $\varsigma^\pm, \bar{\varsigma}^\pm$ being anticommutative, they are also nilpotent, and so only non-negative powers make sense. Moreover, all Taylor series over $\varsigma^\pm, \bar{\varsigma}^\pm$ must terminate with the $\bar{\varsigma}^-\varsigma^+\varsigma^+\varsigma^-$ term. Thus, superfields are always analytic, in fact, multinomials of finite (up to quartic) order over $\varsigma^\pm, \bar{\varsigma}^\pm$.

Superfield components

The component fields are defined as the super-Taylor expansion coefficients

$$\phi \stackrel{\text{def}}{=} \Phi|, \quad \psi_\pm \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_\pm \Phi|, \quad F \stackrel{\text{def}}{=} \frac{1}{2} D_- D_+ \Phi|; \quad (2.5a)$$

$$\bar{\phi} \stackrel{\text{def}}{=} \bar{\Phi}|, \quad \bar{\psi}_\pm \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_\pm \bar{\Phi}|, \quad \bar{F} \stackrel{\text{def}}{=} \frac{1}{2} \bar{D}_+ \bar{D}_- \bar{\Phi}|; \quad (2.5b)$$

are the component fields of the chiral and antichiral spin-0 superfield. Note that both $F, \bar{F}$ change sign with respect to the parity, $\mathbf{P}$; that is, $F, \bar{F}$ are pseudo-scalars if the lowest components, $\phi, \bar{\phi}$, are scalars. Complex conjugation on all components of a chiral superfield may (and herein will) be carried only by the index. That is, we write $\bar{\phi} \rightarrow \phi^\mu$, $\bar{\psi}_\pm \rightarrow \psi_\pm^\mu$, $\bar{F} \rightarrow F^\mu$. Component field content is also specified by $\Phi = (\phi; \psi_\pm; F)$ and $\bar{\Phi} = (\bar{\phi}; \bar{\psi}_\pm; \bar{F})$, leaving the numerical tedium of $\sqrt{2}$ factors and such up to the Reader's preference.

Next,

$$x \stackrel{\text{def}}{=} \Xi|, \quad \xi_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Xi|, \quad \chi_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \Xi|, \quad X \stackrel{\text{def}}{=} \frac{1}{2} D_- \bar{D}_+ \Xi|; \quad (2.5c)$$

$$\bar{x} \stackrel{\text{def}}{=} \bar{\Xi}|, \quad \bar{\xi}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \bar{\Xi}|, \quad \bar{\chi}_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_+ \bar{\Xi}|, \quad \bar{X} \stackrel{\text{def}}{=} \frac{1}{2} D_+ \bar{D}_- \bar{\Xi}|; \quad (2.5d)$$

are the component fields of a twisted chiral and a twisted antichiral spin-0 superfield. The fields $X, \bar{X}$ change sign (are pseudoscalar) with respect to the combined parity and complex conjugation, $\mathbf{CP}$, if $x, \bar{x}$ are scalars with respect to it. Complex conjugation on the components of a twisted chiral superfield may again be carried only by the index, at the expense of labeling the ς^- - and the $\bar{\varsigma}^+$ -components by different characters⁵⁾, as was done in (2.5c,d). Appending indices is then straightforward: $\bar{x} \rightarrow x^{\bar{\alpha}}$, $\bar{\xi}_- \rightarrow \xi_-^{\bar{\alpha}}$, $\bar{\chi}_+ \rightarrow \chi_+^{\bar{\alpha}}$ and $\bar{X} \rightarrow X^{\bar{\alpha}}$. As with the chiral superfields, we may write $\Xi = (x; \xi_-, \chi_+; X)$ and $\bar{\Xi} = (\bar{x}; \bar{\xi}_-, \bar{\chi}_+; \bar{X})$.

Finally, we define

$$\ell \stackrel{\text{def}}{=} \Lambda|, \quad \lambda_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_+ \Lambda|, \quad \lambda_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \Lambda|, \quad L_\mp \stackrel{\text{def}}{=} \frac{1}{4} [D_+, \bar{D}_+] \Lambda|; \quad (2.5e)$$

and

$$r \stackrel{\text{def}}{=} \Upsilon|, \quad \rho_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Upsilon|, \quad \varrho_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \Upsilon|, \quad R_- \stackrel{\text{def}}{=} \frac{1}{4} [D_-, \bar{D}_-] \Upsilon|; \quad (2.5f)$$

⁵⁾ The alternative is to meticulously retain the ‘dot-over’ on spin indices stemming from the $\bar{D}$'s, as was done in Ref. [4]; for some further details, see the appendix A.

are the component fields in the lepton and righton superfields. Note that both $L_{\pm}, R_{\pm}$ change sign under complex conjugation if ℓ, r do not: $L_{\pm}, R_{\pm}$ are imaginary if ℓ, r are real. In general, however, these superfields and their components are complex. Components of the conjugate superfields are defined as hermitian conjugates of (2.5e, f), and so

$$\bar{\ell} \stackrel{\text{def}}{=} \bar{\Lambda}|, \quad \bar{\lambda}_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \bar{\Lambda}|, \quad \bar{\lambda}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_+ \bar{\Lambda}|, \quad \bar{L}_{\pm} \stackrel{\text{def}}{=} \frac{1}{4} [D_+, \bar{D}_+] \bar{\Lambda}|; \quad (2.5\bar{e})$$

and

$$\bar{r} \stackrel{\text{def}}{=} \bar{\Upsilon}|, \quad \bar{\rho}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \bar{\Upsilon}|, \quad \bar{\rho}_+ \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \bar{\Upsilon}|, \quad \bar{R}_{\pm} \stackrel{\text{def}}{=} \frac{1}{4} [D_-, \bar{D}_-] \bar{\Upsilon}|; \quad (2.5\bar{f})$$

Again, we may write $\Lambda = (\ell; \lambda_+, \lambda_-; L_{\pm})$ and $\Upsilon = (r; \rho_-, \rho_+; R_{\pm})$, and $\bar{\Lambda} = (\bar{\ell}; \bar{\lambda}_+, \bar{\lambda}_-; \bar{L}_{\pm})$ and $\bar{\Upsilon} = (\bar{r}; \bar{\rho}_-, \bar{\rho}_+; \bar{R}_{\pm})$.

A remark is perhaps in order. The astute Reader must have wondered about the seemingly unequal definitions of $L_{\pm}, R_{\pm}$ in (2.5e, f), as compared to those of $F, \bar{F}, X, \bar{X}$ in Eqs. (2.5a, b, c, d). Since, e.g., $[D_-, D_+] = 2D_- D_+$, the original definition of F in (2.5a) is merely a minor simplification of $F = \frac{1}{4} [D_-, D_+] \Phi$, and the analogous applies to the definitions (2.5b–d). However, since $[D_+, \bar{D}_+] = 2D_+ \bar{D}_+ - 2i\partial_+$, the definitions of $L_{\pm}, R_{\pm}$ in Eqs. (2.5e, f) imply

$$D_+ \bar{D}_+ \Lambda| = i\partial_+ \Lambda| + \frac{1}{2} [D_+, \bar{D}_+] \Lambda| = i\partial_+ \ell + 2L_+, \quad (2.6a)$$

$$D_- \bar{D}_- \Upsilon| = i\partial_- \Upsilon| + \frac{1}{2} [D_-, \bar{D}_-] \Upsilon| = i\partial_- r + 2R_-. \quad (2.6b)$$

These definitions will produce more symmetry in derivations of Lagrangian densities.

Component diagrammatics

In analyzing the component field content of constrained superfields, the Reader may find it convenient to use the diagram in Fig. 1. A component is obtained by acting on the superfield with one of the operators from the diagram, and then setting $\varsigma^{\pm}=0=\bar{\varsigma}^{\pm}$.




Figure 1. The sequence of multiple superderivatives used in defining component fields. Component fields of conjugate superfields are found using the hermitian conjugates of the operators shown here.

Applying further (super)derivatives merely produces total derivatives of the already obtained component fields. In this sense, the component fields, onto which the operators in the diagram in Fig. 1 project (upon setting $\varsigma^{\pm}=0=\bar{\varsigma}^{\pm}$), are *independent*. The Reader

of a more mathematical persuasion may appreciate the fact that the diagram in Fig. 1 shows the projection operators (with the setting of $\varsigma=0=\bar{\varsigma}$ understood) on the components of $\wedge^*V$, where V is a 4-dimensional vector space spanned by $D_{\pm}, \bar{D}_{\pm}$. The diagram itself then represents the exact sequence

$$1 \rightarrow V \rightarrow \wedge^2 V \rightarrow \wedge^3 V \rightarrow \wedge^4 V . \quad (2.7)$$

Using the diagram in Fig. 1 as a guide, the components of a general, unconstrained complex superfield $\mathbf{A}$ may be listed as

$$\mathbf{A} = \begin{pmatrix} A_{=} \\ \alpha_{-} & A & @_{-} \\ a_{-} & A_{\mp} & \tilde{@}_{-} \\ \alpha_{+} & A_{\pm} & @_{+} \\ a_{+} & A & \tilde{@}_{+} \\ A_{\mp} & & \Delta \end{pmatrix} \quad (2.8)$$

Note that the subscripts on $A_{\mp}, A_{\pm}$ indicate that they were obtained from $[D_{-}, \bar{D}_{+}]\mathbf{A}|$ and $[D_{+}, \bar{D}_{-}]\mathbf{A}|$, respectively. Finally, the numerical coefficients, as in (2.5a), are herein set: $\frac{1}{\sqrt{2}}$ for every superderivative, and $\frac{1}{2}$ for every commutator. Thus, for example,

$$\begin{aligned} a &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_{-} \mathbf{A}| , & A &\stackrel{\text{def}}{=} \frac{1}{2} D_{-} D_{+} \mathbf{A}| , & A_{\pm} &\stackrel{\text{def}}{=} \frac{1}{2} D_{+} \bar{D}_{-} \mathbf{A}| , \\ A &\stackrel{\text{def}}{=} \frac{1}{2} \bar{D}_{+} \bar{D}_{-} \mathbf{A}| , & @_{+} &\stackrel{\text{def}}{=} \frac{1}{4\sqrt{2}} [D_{+}, \bar{D}_{+}] D_{-} \mathbf{A}| , & \text{etc.} \end{aligned} \quad (2.9)$$

The minor simplifying departure in Eqs. (2.5c, d) from this convention, which dictates $X_{\mp}, \bar{X}_{\pm}$ instead of the simpler $X, \bar{X}$, hopefully causes no undue confusion.

Elimination of component fields

Consider, for example, a superfield annihilated by $\bar{D}_{+}$. The component fields obtained by projection after acting with some superderivatives including $\bar{D}_{+}$ either vanish or become total derivatives of ‘lower’ components. All these components are connected by the criss-cross lines in Fig. 1, following to the right of $\bar{D}_{+}$; they are shown in the shaded area of the diagram in Fig 2.




Figure 2. The (independent) component fields eliminated by the superconstraint $\bar{D}_{+}\Psi=0$ are obtained using the derivatives in the shaded area. Thus, the superfield Ψ , satisfying $\bar{D}_{+}\Psi=0$ retains the independent components obtained by projection with the unshaded superderivatives.

The superconstraint of being annihilated by $\bar{D}_-$ on the other hand eliminates the (independent) component fields obtained by projection with the derivatives in the shaded area of the diagram in Fig. 3.




Figure 3. The (independent) component fields eliminated by the superconstraint $\bar{D}_-\Psi=0$ are obtained using the derivatives in the shaded area.

The chiral superfield, Φ , satisfies both $\bar{D}_+\Phi=0$ and $\bar{D}_-\Phi=0$, so the (independent) component fields eliminated by both superconstraints are in the *union* of the shaded areas of Figs. 2 and 3; this is shown in Fig. 4.




Figure 4. The (independent) component fields eliminated by the superconstraint pair $\bar{D}_-\Phi=0=\bar{D}_+\Phi$ are obtained using the derivatives in the shaded area; their overlap is shaded darker. Unshaded remain the operators which project on the independent components of the chiral superfield, Φ : see Eqs. (2.5a).

Note that there is an overlap of the two shaded areas: the derivatives projecting onto the component fields that are eliminated both by $\bar{D}_+\Phi=0$ and by $\bar{D}_-\Phi=0$. Presently, this is of no concern, but will turn up again later.

Clearly, the component field content of all constrained superfields (2.1) and (2.3) can be treated in much the same way. Unfortunately, while this does give a quick indication of the *independent* component field content, it does not specify the dependences. For example, such diagrams *do* indicate that the right-hand side of Eq. (2.6a) is a linear combination of $L_\pm$ and a derivative (clearly, $\partial_\pm$) of a component field two columns to the left in the diagram (clearly, ℓ). Such diagrams, *by themselves*, do not however specify the numerical constants of the linear combination, here ‘2’ and ‘i’, respectively. These details, while not difficult to guess, should be verified by direct calculation.

Unidexterity superconstraints

We have noted before that the pair of superconstraints (2.1e) *includes* the spacetime constraint, $\partial_=\Lambda = 0$, and that (2.1f) similarly *includes* $\partial_+\Upsilon = 0$. The quick argument based on

$$\left. \begin{array}{l} D_-\Lambda = 0 \\ \bar{D}_-\Lambda = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} \bar{D}_-D_-\Lambda = 0 \\ D_-\bar{D}_-\Lambda = 0 \end{array} \right\} \Rightarrow \{D_-, \bar{D}_-\}\Lambda = 2i\partial_=\Lambda = 0, \quad (2.10)$$

is somewhat misleading in suggesting that $(\partial_=\Lambda)=0$ is somehow a *consequence* of (2.1e), and so a secondary constraint; a clarification may thus be welcome. To that end, let us study the content of Eqs. (2.1e) on the component fields. Start with an unconstrained (general) superfield, $\mathring{\Lambda}$, and define ‘ $\mathring{\cdot}$ ’ to mean setting $\varsigma^-=0=\bar{\varsigma}^-$, but leaving $\varsigma^+, \bar{\varsigma}^+$ intact. Then, we define the (*still* $\varsigma^+, \bar{\varsigma}^+$ -dependent!) $\varsigma^-, \bar{\varsigma}^-$ -component (0,2)-superfields:

$$\Lambda \stackrel{\text{def}}{=} \mathring{\Lambda} \rfloor, \quad \mathring{\lambda}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \mathring{\Lambda} \rfloor, \quad \mathring{\bar{\lambda}}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \mathring{\Lambda} \rfloor, \quad \mathring{L}_- \stackrel{\text{def}}{=} \frac{1}{4} [D_-, \bar{D}_-] \mathring{\Lambda} \rfloor, \quad (2.11)$$

or $\mathring{\Lambda} = (\Lambda; \mathring{\lambda}_-, \mathring{\bar{\lambda}}_-; \mathring{L}_-)$.

Now project on the $\varsigma^-, \bar{\varsigma}^-$ -component constraints (components of $D_- \mathring{\Lambda} \stackrel{!}{=} 0$):

$$0 \stackrel{!}{=} (D_- \mathring{\Lambda}) \rfloor = \sqrt{2} \mathring{\lambda}_-, \quad (2.12a)$$

$$0 \stackrel{!}{=} \frac{1}{\sqrt{2}} \bar{D}_- (D_- \mathring{\Lambda}) \rfloor = \frac{i}{\sqrt{2}} (\partial_=\Lambda) + \sqrt{2} \mathring{L}_-, \quad (2.12b)$$

$$0 \stackrel{!}{=} \frac{1}{4} [D_-, \bar{D}_-] (D_- \mathring{\Lambda}) \rfloor = \frac{i}{\sqrt{2}} (\partial_=\mathring{\lambda}_-), \quad (2.12c)$$

and of $\bar{D}_- \mathring{\Lambda} \stackrel{!}{=} 0$:

$$0 \stackrel{!}{=} (\bar{D}_- \mathring{\Lambda}) \rfloor = \sqrt{2} \mathring{\bar{\lambda}}_-, \quad (2.12d)$$

$$0 \stackrel{!}{=} \frac{1}{\sqrt{2}} D_- (\bar{D}_- \mathring{\Lambda}) \rfloor = \frac{i}{\sqrt{2}} (\partial_=\Lambda) - \sqrt{2} \mathring{L}_-, \quad (2.12e)$$

$$0 \stackrel{!}{=} \frac{1}{4} [D_-, \bar{D}_-] (\bar{D}_- \mathring{\Lambda}) \rfloor = \frac{i}{\sqrt{2}} (\partial_=\mathring{\bar{\lambda}}_-), \quad (2.12f)$$

where $\stackrel{!}{=}$ is the equality of the enforced (super)constraint. We have used only the *operatorial* identity

$$\{D_-, \bar{D}_-\} = 0, \quad \Rightarrow \quad [D_-, \bar{D}_-] D_- \equiv \{D_-, \bar{D}_-\} D_- , \quad (2.13)$$

and its conjugate.

Clearly, the sum of Eqs.(2.12b) and (2.12e) equals the spacetime constraint, $\partial_=\Lambda \stackrel{!}{=} 0$, for *all* of the haploid (0,2)-superfield Λ . Their difference produces $\mathring{L}_- \stackrel{!}{=} 0$ —which is in point of fact also *contained* both in (2.12a) and in (2.12d). To summarize then, the superconstraints (2.1e) *contain* the (simple) constraints:

$$(\partial_=\Lambda) \stackrel{!}{=} 0, \quad \text{and} \quad \mathring{\lambda}_-, \mathring{\bar{\lambda}}_-, \mathring{L}_- \stackrel{!}{=} 0. \quad (2.14)$$

That is, the vanishing of three quarters ($\mathring{\lambda}_-, \mathring{\bar{\lambda}}_-, \mathring{L}_-$) of the unconstrained superfield $\mathring{\Lambda}$ and the vanishing of the right-moving modes of all the component fields in the haploid superfield Λ are all *contained* in Eqs. (2.1e); their content is *identical* to that of Eqs. (2.14)!

Thus, the various stages (columns) of the ‘derivation’ (2.10) are *consistent*, but also unnecessary. Eqs. (2.14) merely expand the original superdifferential system (2.1e), which contains all the (super)constraints; there are no further, induced, (super)constraints to be derived in the Lagrangian formalism used here. The strategy in the Hamiltonian (Dirac, BRST, *etc.*) formalism will depend crucially on the conjugate momenta, and so on the choice of the Lagrangian density. This again is beyond our present scope, as we seek to present the (reasonably) general case rather than belabor the details—and consequent restrictions—of specific models.

Quartoid superfield components

The component fields of the quartoid superfields (2.3) are easily obtained, by truncation, from those of the haploid ones (2.5):

$$\phi_L \stackrel{\text{def}}{=} \Phi_L|, \quad \psi_{L+} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_+ \Phi_L|; \quad (2.15a)$$

$$\bar{\phi}_L \stackrel{\text{def}}{=} \bar{\Phi}_L|, \quad \bar{\psi}_{L+} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \bar{\Phi}_L|; \quad (2.15b)$$

$$\phi_R \stackrel{\text{def}}{=} \Phi_R|, \quad \psi_{R-} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Phi_R|; \quad (2.15c)$$

$$\bar{\phi}_R \stackrel{\text{def}}{=} \bar{\Phi}_R|, \quad \bar{\psi}_{R-} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \bar{\Phi}_R|. \quad (2.15d)$$

All the superfields defined in Eqs. (2.1) and (2.3) are spin-0 superfields, in that their lowest components (2.5) are all spin-0 fields. For these to be standard physical (pseudo)scalar fields, we fix their canonical dimension to be the standard 0. Thence, the canonical dimensions of all spinors in (2.5) is the standard $+\frac{1}{2}$.

2.4. *Spinning superfields and superdifferential relations*

The Eqs. (2.1) define *scalar* superfields, *i.e.*, the lowest component fields of $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}, \Lambda, \Upsilon$ are scalar fields. It is easy to endow the whole superfield with (overall) additional spin, say $\Phi \rightarrow \Phi^+$, so that the lowest component $\Phi^+| = \phi^+$ has spin $+\frac{1}{2}$ (see appendix A). Next, it is also possible to relate the (anti)chiral and the twisted (anti)chiral superfields via a superdifferential relation. For example, let Φ^+ be a spin-up (overall spin $+\frac{1}{2}$) chiral superfield; then $\Xi' \stackrel{\text{def}}{=} (D_+ \Phi^+)$ is a scalar (overall spin 0) twisted chiral superfield, since

$$D_+(D_+ \Phi^+) = 0 = \bar{D}_-(D_+ \Phi^+). \quad (2.16)$$

In particular, note that although $\bar{D}_+ \Phi^+ = 0$,

$$\bar{D}_+(D_+ \Phi^+) = (2i\partial_{\neq} - D_+ \bar{D}_+) \Phi^+ = 2i\partial_{\neq} \Phi^+ \neq 0. \quad (2.17)$$

This then implies that there will exist (differential) relations among the component fields

of Φ^+ and $\Xi' \stackrel{\text{def}}{=} (D_+ \Phi^+)$, as follows:

$$\begin{aligned}
x' &\stackrel{\text{def}}{=} \Xi'| = (D_+ \Phi^+)| = \sqrt{2} \psi_+^+ , \\
\xi'_- &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Xi'| = \frac{1}{\sqrt{2}} (D_- D_+ \Phi^+)| = \sqrt{2} F^+ , \\
\xi'_+ &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \Xi'| = \frac{1}{\sqrt{2}} (\bar{D}_+ D_+ \Phi^+)| = \sqrt{2} i (\partial_{\mp} \phi^+) , \\
X' &\stackrel{\text{def}}{=} \frac{1}{2} D_- \bar{D}_+ \Xi'| = \frac{1}{2} (D_- \bar{D}_+ D_+ \Phi^+)| = \sqrt{2} i (\partial_{\mp} \psi_-^+) .
\end{aligned}$$

Also, since the canonical (scaling) dimension of $D_{\pm}$ and $\bar{D}_{\pm}$ is $[D_{\pm}] = [\bar{D}_{\pm}] = +\frac{1}{2}$, the component fields in $\Xi' \stackrel{\text{def}}{=} D_+ \Phi^+$ will have their canonical dimensions shifted by $+\frac{1}{2}$ compared to those in Φ^+ . For example, the lowest component of Φ^+ is the spinor ϕ^+ and so should (in 2-dimensional spacetime), have $[\phi^+] = \frac{1}{2}$; then $[\psi_{\pm}^+] = 1$ and $[F^+] = \frac{3}{2}$. Thus, $[x'] = 1$, $[\xi'_{\pm}] = \frac{3}{2}$, and $[X'] = 2$ —none of which are the ‘correct’ canonical dimensions of physical (propagating) fields in 2-dimensional spacetime. If a mass/energy parameter, m , is available in the model, the rescaled superfield $\tilde{\Xi} \stackrel{\text{def}}{=} \frac{1}{m} (D_+ \Phi^+)$ has component fields of ‘correct’ (physical) canonical dimensions.

Alternatively, with Ξ a scalar (overall spin-0) twisted chiral superfield, $\Phi_+ \stackrel{\text{def}}{=} (\bar{D}_+ \Xi)$ is a spin $-\frac{1}{2}$ chiral superfield since

$$\bar{D}_+ (\bar{D}_+ \Xi) = 0 = \bar{D}_- (\bar{D}_+ \Xi) . \quad (2.18)$$

Again, the component fields of Ξ and $\Phi_+ \stackrel{\text{def}}{=} (\bar{D}_+ \Xi)$ are related as follows:

$$\begin{aligned}
\phi_+ &\stackrel{\text{def}}{=} \Phi_+| = (\bar{D}_+ \Xi)| = \sqrt{2} \chi_+ , \\
\psi &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Phi_+| = \frac{1}{\sqrt{2}} (D_- \bar{D}_+ \Xi)| = \sqrt{2} X , \\
\psi_{\mp} &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_+ \Phi_+| = \frac{1}{\sqrt{2}} (D_+ \bar{D}_+ \Xi)| = \sqrt{2} i (\partial_{\mp} x) , \\
F_+ &\stackrel{\text{def}}{=} \frac{1}{2} D_- D_+ \Phi_+| = \frac{1}{2} (D_- D_+ \bar{D}_+ \Xi)| = \sqrt{2} i (\partial_{\mp} \xi_-) .
\end{aligned}$$

Again, the canonical (scaling) dimensions of the component fields of $\Phi_+ \stackrel{\text{def}}{=} (\bar{D}_+ \Xi)$ will each have their canonical dimensions shifted by $+\frac{1}{2}$ compared to those in Ξ . So, if all the component fields of Ξ have ‘correct’ canonical dimensions, not all of those of Φ_+ do. Whilst $[\phi_+] = [\chi_+] = \frac{1}{2}$ is the correct canonical dimension for a dynamical spinor, and $[\psi_{\mp}] = [\partial_{\mp} x] = 1$ and $[\psi] = [F] = 1$ are correct for a component of a vector and an auxiliary field, respectively, $[F_+] = [\partial_{\mp} \xi_-] = \frac{3}{2}$ is too high for a dynamical spinor.

This suggests that superderivatives of superfields generally have fewer dynamical component fields than the original superfields. Furthermore, the unphysical canonical dimensions of the component fields of superderivative superfields allow for higher derivative terms in the action. Throughout this article and with one noted exception below, we will include such superderivative superfields only if this does not lead to higher derivative terms.

Note that no superderivative of either of $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}$ satisfies the unidexterous superconstraints (2.1e, f). Thus, it is not possible to construct lefton or righton superfields from the (twisted) (anti)chiral superfields $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}$.

Conversely however, superderivatives of leftons and rightons do satisfy some of the (twisted) (anti)chiral conditions. In fact, these are annihilated by *three* superderivatives and are *quartoid*, no longer merely haploid, superfields. Thus:

- $(\bar{D}_+\Lambda)$ is a chiral lefton (2.3a), and $(D_+\Lambda)$ an antichiral lefton (2.3b),
- $(\bar{D}_-\Upsilon)$ is a chiral righton (2.3c), and $(D_-\Upsilon)$ an antichiral righton (2.3d).

2.5. Second-order constrained superfields

The hitherto discussed and thoroughly familiar (anti)chiral superfields (2.1a, b), their somewhat less well known twisted counterparts (2.1c, d), and the unidexterous fields (2.1e, f) and (2.3), are all defined by satisfying a system of *first order* superconstraints. Clearly, there is more.

The obvious generalization of the (quite simple) first-order superconstraints (2.1) and (2.3) are the (equally simple) *second-order* superconstraints:

$$1. \text{ NM-Chiral : } \quad \bar{D}_+\bar{D}_-\Theta = 0 , \quad (2.19a)$$

$$2. \text{ NM-Antichiral : } \quad D_-D_+\bar{\Theta} = 0 , \quad (2.19b)$$

$$3. \text{ NM-Twisted-chiral : } \quad D_+\bar{D}_-\Pi = 0 , \quad (2.19c)$$

$$4. \text{ NM-Twisted-antichiral : } \quad D_-\bar{D}_+\bar{\Pi} = 0 , \quad (2.19d)$$

$$5. \text{ NM-(Almost)-Lefton : } \quad [D_-, \bar{D}_-]\mathbf{A} = 0 , \quad (2.19e)$$

$$6. \text{ NM-(Almost)-Righton : } \quad [D_+, \bar{D}_+]\mathbf{U} = 0 . \quad (2.19f)$$

These are called ‘linear’, or as better befits the extendedness of their defining superconstraints in comparison with (2.1), ‘non-minimal’ [12,7,13]. Note that the ‘minimal’ haploid superfields (2.1) obey the second-order superconstraints (2.19), but that the non-minimal ones (2.19) do not obey the first-order superconstraints (2.1).

Again, the latter two of the superconstraints (2.19) are set apart from the former four, although the divide is not as sharp as between the ‘minimal’ unidexterous superfields (2.1e, f) from the ambidexterous ones (2.1a–d). To see this, let us examine their component field content.

Ambidexterous non-minimal superfields

The component field content is straightforward to obtain, only this time there are more component fields than in Eqs. (2.5). For example,

$$\begin{aligned} t &\stackrel{\text{def}}{=} \Theta| , \quad \theta_{\pm} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}}D_{\pm}\Theta| , \quad \vartheta_{\pm} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}}\bar{D}_{\pm}\Theta| , \\ T &\stackrel{\text{def}}{=} \frac{1}{2}D_-D_+\Theta| , \quad T_{\mp} \stackrel{\text{def}}{=} \frac{1}{2}D_-\bar{D}_{\mp}\Theta| , \quad T_{\pm} \stackrel{\text{def}}{=} \frac{1}{2}D_+\bar{D}_{\pm}\Theta| , \\ T_{\pm} &\stackrel{\text{def}}{=} \frac{1}{4}[D_-, \bar{D}_{\pm}]\Theta| , \quad T_{\mp} \stackrel{\text{def}}{=} \frac{1}{4}[D_+, \bar{D}_{\mp}]\Theta| , \\ \tau_{-} &\stackrel{\text{def}}{=} \frac{1}{4\sqrt{2}}[D_-, \bar{D}_{-}]D_+\Theta| , \quad \tau_{+} \stackrel{\text{def}}{=} \frac{1}{4\sqrt{2}}[D_+, \bar{D}_{+}]D_-\Theta| . \end{aligned} \quad (2.20a)$$

Note that the $\pm$ and $\mp$ subscripts on $T_{\pm}, T_{\mp}$ indicate *single* components, whereas the $\pm$ on the fermions $\theta_{\pm}, \vartheta_{\pm}, \tau_{\pm}$ indicates a *choice* of spin $+\frac{1}{2}$ and $-\frac{1}{2}$. Similarly,

$$\begin{aligned}
p &\stackrel{\text{def}}{=} \Pi|, \quad \pi_{\pm} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_{\pm} \Pi|, \quad \varpi_{\pm} \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_{\pm} \Pi|, \\
P &\stackrel{\text{def}}{=} \frac{1}{2} D_- D_+ \Pi|, \quad P_{\mp} \stackrel{\text{def}}{=} \frac{1}{2} D_- \bar{D}_+ \Pi|, \quad \mathbb{P} \stackrel{\text{def}}{=} \frac{1}{2} \bar{D}_+ \bar{D}_- \Pi|, \\
P_{\pm} &\stackrel{\text{def}}{=} \frac{1}{4} [D_-, \bar{D}_-] \Pi|, \quad P_{\mp} \stackrel{\text{def}}{=} \frac{1}{4} [D_+, \bar{D}_+] \Pi|, \\
\tilde{\varphi}_{-} &\stackrel{\text{def}}{=} \frac{1}{4\sqrt{2}} [D_-, \bar{D}_-] \bar{D}_+ \Pi|, \quad \varphi_{+} \stackrel{\text{def}}{=} \frac{1}{4\sqrt{2}} [D_+, \bar{D}_+] D_- \Pi|.
\end{aligned} \tag{2.20b}$$

The component fields of $\bar{\Theta}$ and $\bar{\Pi}$ are then obtained by hermitian conjugation.




Figure 5. The (independent) component fields eliminated by the superconstraint $\bar{D}_+ \bar{D}_- \Theta = \frac{1}{2} [\bar{D}_+, \bar{D}_-] \Theta = 0$ are obtained using the derivatives in the shaded area. The unshaded derivatives project on the independent component fields of Θ , as defined by Eqs. (2.20.)

Note that the shaded area of Fig. 5 coincides with the darker shaded area of Fig. 4. Indeed, trivially so: the former covers the derivatives defining the component fields eliminated by projection with $\bar{D}_+ \bar{D}_-$, the latter — with either of $\bar{D}_+, \bar{D}_-$.

Comparing the component field content of corresponding superfields, the non-minimal superfields (2.19a–d) contain three times as many component fields as do the ‘minimal’ ones (2.1a–d). As it turns out, however, with rather typical Lagrangian densities for such superfields (even when interacting with some of the others), the equations of motion for the ‘extra’ component fields turn out to be algebraic (see, *e.g.*, Ref.[7], p.200). This allows an easy elimination of all the ‘extra’ component fields, and the remaining component field content of the non-minimal superfields (2.19a–d) turns out to be identical to their ‘minimal’ counterparts (2.1a–d). To indicate this, we may list the component field content as

$$\Theta = (t; \theta_{\pm}; T : \bar{\vartheta}_{\pm}; T_{\mp}, T_{\pm}, T_-, T_{\mp}; \tau_{\pm}), \tag{2.21a}$$

$$\bar{\Theta} = (\bar{t}; \bar{\theta}_{\pm}; \bar{T} : \vartheta_{\pm}; \bar{T}_{\mp}, \bar{T}_{\pm}, \bar{T}_-, \bar{T}_{\mp}; \bar{\tau}_{\pm}), \tag{2.21b}$$

$$\Pi = (p; \pi_-, \varpi_+; P : \varpi_-, \pi_+; P_{\mp}, \mathbb{P}, P_-, P_{\mp}; \tilde{\varphi}_-, \varphi_+), \tag{2.21c}$$

$$\bar{\Pi} = (\bar{p}; \bar{\pi}_-, \bar{\varpi}_+; \bar{P} : \bar{\varpi}_-, \bar{\pi}_+; \bar{\mathbb{P}}, \bar{P}_{\pm}, \bar{P}_-, \bar{P}_{\mp}; \tilde{\bar{\varphi}}_-, \bar{\varphi}_+), \tag{2.21d}$$

where the colon separates off the ‘extra’ component fields by which non-minimal superfields differ from their ‘minimal’ counterparts.

Thus, the physical component field content warrants the adjective ‘haploid’ also for the non-minimal superfields, as they depend on the ‘other half’ of the four supercoordinates, $\varsigma^\pm, \bar{\varsigma}^\pm$ only by the inclusion of auxiliary component fields.

Furthermore, the non-minimal superfields (2.19) are *dual* to their first-order constrained counterparts (2.1) (see, *e.g.*, Ref. [7], p.200). This tempts a straightforward dismissal of the non-minimal superfields. However, while the use of these non-minimal superfields *instead* of the ‘minimal’ ones seems to merely harbor unnecessary toil, there do exist distinct benefits of using the ‘minimal’ and the non-minimal superfields *together* [13]. We will therefore include them in the subsequent analysis, where appropriate.




Figure 6. The darkest shade indicates the derivatives defining the (independent) component fields eliminated by the superconstraint $\bar{D}_+ \bar{D}_- \Theta = 0$. The medium and lightest shade indicate the two groups of derivatives which jointly define the ‘extra’ component fields. Compare this with Fig. 4.

The above results are displayed diagrammatically in Fig. 6. Note that the derivatives in the lightest and medium shaded areas separately each have the structure of the derivatives defining the (independent) component fields of a ‘minimal’ chiral superfield. This motivated the realization that the definition (2.19a) is invariant with respect to the (rather peculiar) inhomogeneous gauge transformation

$$\delta\Theta = \bar{\varsigma}^- \Phi_- + \bar{\varsigma}^+ \Phi_+, \quad \text{where} \quad \bar{D}_\pm \Phi_\pm = 0 = \bar{D}_\pm \Phi_\mp, \quad (2.22a)$$

parametrized by a spin $\pm\frac{1}{2}$ pair of chiral superfields, $\Phi_\pm$. Indeed, the (independent) components of $\bar{\varsigma}^- \Phi_-$ and $\bar{\varsigma}^+ \Phi_+$ are projected precisely by those derivatives which are located in the lightly- and medium-shaded area of Fig. 6—the ‘extra’ component fields of Θ . The component fields in $\Phi_\pm$ then can be used to gauge away the ‘extra’ component fields in Θ . In other words,

$$\Theta \bmod (\bar{\varsigma}^- \Phi_- + \bar{\varsigma}^+ \Phi_+) \simeq \Phi. \quad (2.23)$$

By the same token, the definitions (2.19b–d) are, respectively, invariant with respect to the following inhomogeneous gauge transformations:

$$\delta\bar{\Theta} = \varsigma^- \bar{\Phi}_- + \varsigma^+ \bar{\Phi}_+, \quad \text{where} \quad D_- \bar{\Phi}_\pm = 0 = D_+ \bar{\Phi}_\pm, \quad (2.22b)$$

$$\delta\Pi = \varsigma^- \Xi_- + \varsigma^+ \Xi_+, \quad \text{where} \quad \bar{D}_- \Xi_\pm = 0 = D_+ \Xi_\pm, \quad (2.22c)$$

$$\delta\bar{\Pi} = \bar{\varsigma}^- \bar{\Xi}_- + \bar{\varsigma}^+ \bar{\Xi}_+, \quad \text{where} \quad D_- \bar{\Xi}_\pm = 0 = \bar{D}_+ \bar{\Xi}_\pm. \quad (2.22d)$$

Of course, most Lagrangian densities will explicitly violate this gauge symmetry (2.22), and a possible use of this gauge symmetry remains an open question for a later study.

(Almost) Unidexterous non-minimal superfields

Before we specify the component field content of the non-minimal superfields (2.19e, f), let us examine the content of the second order superdifferential constraints (2.19e, f). To this end, consider a previously unconstrained general (2,2)-superfield, $\mathring{\mathbf{A}}$ with (*still* ς^+ , $\bar{\varsigma}^+$ -dependent!) component (0,2)-superfield content $\mathring{\mathbf{A}} = (\mathbf{A}; \mathring{\alpha}_-, \mathring{a}_-; \mathring{A}_=)$, and project on the components of $[D_-, \bar{D}_-]\mathring{\mathbf{A}}$:

$$0 \stackrel{!}{=} ([D_-, \bar{D}_-]\mathring{\mathbf{A}}) \quad = \mathring{A}_=, \quad (2.24a)$$

$$0 \stackrel{!}{=} (\frac{1}{\sqrt{2}}D_-[D_-, \bar{D}_-]\mathring{\mathbf{A}}) \quad = -2i(\partial_=\mathring{\alpha}_-), \quad (2.24b)$$

$$0 \stackrel{!}{=} (\frac{1}{\sqrt{2}}\bar{D}_-[D_-, \bar{D}_-]\mathring{\mathbf{A}}) \quad = +2i(\partial_=\mathring{a}_-), \quad (2.24c)$$

$$0 \stackrel{!}{=} (\frac{1}{2}[D_-, \bar{D}_-][D_-, \bar{D}_-]\mathring{\mathbf{A}}) \quad = -(\partial_=\mathring{\mathbf{A}}). \quad (2.24d)$$

The component fields are easily defined in analogy with Eqs. (2.20), and we denote them $\mathbf{A} = (a; \alpha_+, a_+; A_{\mp} : \alpha_-, a_-; A_=: A, A_{\mp}, A_{\pm}, \mathbf{A}; @_+, \tilde{@}_+)$.

Just as with the ambidexterous non-minimal superfields, the defining superconstraint (2.19e) is again invariant with respect to the inhomogeneous gauge transformation

$$\delta \mathbf{A} = \varsigma^- \Lambda_-^1 + \bar{\varsigma}^- \Lambda_-^2, \quad \text{where} \quad D_- \Lambda_-^i = 0 = \bar{D}_- \Lambda_-^i, \quad i = 1, 2, \quad (2.22e)$$

which precisely suffices to eliminate the ‘extra’ component fields, $\alpha_-, a_-, A_=: A, A_{\mp}, A_{\pm}, \mathbf{A}, @_+, \tilde{@}_+$, all of which are contained in $\mathring{\alpha}_-$ and $\mathring{a}_-$. The remaining component fields are $(a; \alpha_+, a_+; A_{\mp})$, and are contained in $\mathbf{A}$ but neither in $\mathring{\alpha}_-$ nor in $\mathring{a}_-$. This has the peculiar consequence that the component fields $(a; \alpha_+, a_+; A_{\mp})$ obey the superconstraint (2.24d), but not (2.24b, c). That is, the component fields $(a; \alpha_+, a_+; A_{\mp})$ are not purely left-movers, but are linear functions of $\sigma^=$ and arbitrary functions of $\sigma^{\pm}$.

The analogous analysis shows that the non-minimal righton has the following component fields $\mathbf{U} = (u; \nu_-, \nu_-; U_=: \nu_+, \nu_+; U, U_{\mp}, U_{\pm}, \mathbb{U}; \mu_-, \tilde{\mu}_-)$. Its definition is invariant with respect to the gauge transformation

$$\delta \mathbf{U} = \varsigma^+ \Upsilon_+^1 + \bar{\varsigma}^+ \Upsilon_+^2, \quad \text{where} \quad D_+ \Upsilon_+^i = 0 = \bar{D}_+ \Upsilon_+^i, \quad i = 1, 2, \quad (2.22f)$$

which eliminates the ‘extra’ component fields, $\nu_+, \nu_+, U, U_{\mp}, U_{\pm}, \mathbb{U}, \mu_-, \tilde{\mu}_-$, contained in $\mathring{\rho}_+$ and $\mathring{\varrho}_+$. The remaining component fields are $(u; \nu_-, \nu_-; U_=)$, and are linear functions of $\sigma^{\pm}$ and arbitrary functions of $\sigma^=$.

Finally, we recall that an analytic function of any one type of ‘minimal’ haploid (quartoid) superfields (2.1) and (2.3) is again a haploid (quartoid) superfield, and of the same type. For example, any analytic function $f(\Phi)$ of chiral superfields, Φ^{μ} , is again chiral: $f(\Phi)$ satisfies the same superconstraints (2.1a). In contradistinction, only linear combinations of non-minimal superfields of the same type (2.19a–f) satisfy the same superconstraint. No *non-linear* function of any non-minimal superfields satisfies any one of the non-minimal superconstraints (2.19a–f)—they are all general, unconstrained superfields.

Minimization of non-minimal superfields

As observed in the literature [4], superderivatives of non-minimal superfields may become minimal. For example, consider the non-minimal chiral superfield (2.19a), Θ , satisfying $\bar{D}_+ \bar{D}_- \Theta = 0$. Then both $(\bar{D}_+ \Theta)$ and $(\bar{D}_- \Theta)$ are minimal chiral superfields; indeed,

$$\begin{aligned} \bar{D}_+(\bar{D}_+ \Theta) &\equiv 0, & \bar{D}_-(\bar{D}_+ \Theta) &= 0; \\ \bar{D}_-(\bar{D}_- \Theta) &\equiv 0, & \bar{D}_+(\bar{D}_- \Theta) &= 0. \end{aligned} \quad (2.25)$$

However, $(D_+ \Theta)$ and $(D_- \Theta)$ are not even non-minimal haploid superfields: they are annihilated by only one of the four superderivatives.

In this way, every non-minimal ambidexterous superfield (2.19a)-(2.19d) provides two minimal haploid superfields (2.1a-d) of the corresponding kind:

$$(\bar{D}_+ \Theta), (\bar{D}_- \Theta) \quad \text{are chiral (2.1a)}, \quad (2.26a)$$

$$(D_+ \bar{\Theta}), (D_- \bar{\Theta}) \quad \text{are antichiral (2.1b)}, \quad (2.26b)$$

$$(D_+ \Pi), (\bar{D}_- \Pi) \quad \text{are twisted-chiral (2.1c)}, \quad (2.26c)$$

$$(\bar{D}_+ \bar{\Pi}), (D_- \bar{\Pi}) \quad \text{are twisted-antichiral (2.1d)}. \quad (2.26d)$$

The same is not true for the unidexterous superfields. Consider the non-minimal lefton, $\mathbf{A}$, satisfying $[D_-, \bar{D}_-] \mathbf{A} = 0$. Then,

$$\begin{aligned} \bar{D}_-(\bar{D}_- \mathbf{A}) &\equiv 0, \quad \text{but} \quad D_-(\bar{D}_- \mathbf{A}) = i(\partial_- \mathbf{A}) \neq 0; \\ D_-(D_- \mathbf{A}) &\equiv 0, \quad \text{but} \quad \bar{D}_-(D_- \mathbf{A}) = i(\partial_- \mathbf{A}) \neq 0. \end{aligned} \quad (2.27)$$

However, there are analogous relations between unidexterous haploid and quartoid superfields. It is easy to check that

$$(\bar{D}_+ \Lambda) \quad \text{is a chiral lefton (2.3a)}, \quad (2.28a)$$

$$(D_+ \Lambda) \quad \text{is an antichiral lefton (2.3b)}, \quad (2.28b)$$

$$(\bar{D}_- \Upsilon) \quad \text{is a chiral righton (2.3c)}, \quad (2.28c)$$

$$(D_- \Upsilon) \quad \text{is an antichiral righton (2.3d)}. \quad (2.28d)$$

Effectively minimal haploid superfields

To summarize, the following (superderivative) superfields satisfy the specified haploid and quartoid superconstraints:

Chiral, obeying Eq. (2.1a) :

$$\begin{aligned} &\Phi; (\bar{D}_+ \Xi), (\bar{D}_- \Xi), (\bar{D}_+ \Lambda), (\bar{D}_- \Upsilon); (\bar{D}_+ \Theta), (\bar{D}_- \Theta); \\ &(\bar{D}_+ \bar{D}_- \bar{\Phi}), (\bar{D}_+ D_+ \Lambda), (\bar{D}_- D_- \Upsilon), (\bar{D}_+ D_+ \Pi), (\bar{D}_- D_- \bar{\Pi}); \end{aligned} \quad (2.29a)$$

Antichiral, obeying Eq. (2.1b) :

$$\begin{aligned} & \bar{\Phi}; (D_- \Xi), (D_+ \bar{\Xi}), (D_+ \Lambda), (D_- \Upsilon); (D_+ \bar{\Theta}), (D_- \bar{\Theta}); \\ & (D_- D_+ \Phi), (D_+ \bar{D}_+ \Lambda), (D_- \bar{D}_- \Upsilon), (D_- \bar{D}_- \Pi), (D_+ \bar{D}_+ \bar{\Pi}); \end{aligned} \quad (2.29b)$$

Twisted-chiral, obeying Eq. (2.1c) :

$$\begin{aligned} & \Xi; (D_+ \Phi), (\bar{D}_- \bar{\Phi}), (D_+ \Lambda), (\bar{D}_- \Upsilon); (D_+ \Pi), (\bar{D}_- \Pi); \\ & (D_+ \bar{D}_- \bar{\Phi}), (D_+ \bar{D}_+ \Lambda), (\bar{D}_- D_- \Upsilon), (D_+ \bar{D}_+ \Theta), (\bar{D}_- D_- \bar{\Theta}); \end{aligned} \quad (2.29c)$$

Twisted-antichiral, obeying Eq. (2.1d) :

$$\begin{aligned} & \bar{\Xi}; (\bar{D}_+ \bar{\Phi}), (D_- \Phi), (\bar{D}_+ \Lambda), (D_- \Upsilon); (\bar{D}_+ \bar{\Pi}), (D_- \bar{\Pi}); \\ & (D_- \bar{D}_+ \Xi), (\bar{D}_+ D_+ \Lambda), (D_- \bar{D}_- \Upsilon), (D_- \bar{D}_- \Theta), (\bar{D}_+ D_+ \bar{\Theta}); \end{aligned} \quad (2.29d)$$

Leftons, obeying Eq. (2.1e) :

$$\Lambda; (D_+ \Lambda), (\bar{D}_+ \Lambda); (\bar{D}_+ D_+ \Lambda), (D_+ \bar{D}_+ \Lambda); \quad (2.29e)$$

Rightons, obeying Eq. (2.1f) :

$$\Upsilon; (D_- \Upsilon), (\bar{D}_- \Upsilon); (D_- \bar{D}_- \Upsilon), (\bar{D}_- D_- \Upsilon); \quad (2.29f)$$

and

$$\text{Chiral leftons, obeying Eq. (2.3a) :} \quad \Phi_L; (\bar{D}_+ \Lambda); (\bar{D}_+ D_+ \Lambda); \quad (2.29g)$$

$$\text{Antichiral leftons, obeying Eq. (2.3b) :} \quad \bar{\Phi}_L; (D_+ \Lambda); (D_+ \bar{D}_+ \Lambda); \quad (2.29h)$$

$$\text{Chiral rightons, obeying Eq. (2.3c) :} \quad \Phi_R; (\bar{D}_- \Upsilon); (\bar{D}_- D_- \Upsilon); \quad (2.29i)$$

$$\text{Antichiral rightons, obeying Eq. (2.3d) :} \quad \bar{\Phi}_R; (D_- \Upsilon); (D_- \bar{D}_- \Upsilon). \quad (2.29j)$$

The order of derivatives has been terminated at two only so as to prohibit higher derivative and/or negative-dimension coefficient terms in the Lagrangian density.

Finally, as it will be useful when constructing Lagrangian densities, we list (superderivative) superfields which are annihilated by at least one of the superderivatives. Any analytic function of these is then also annihilated by at least that one superderivative, and forms the *kernel* of that superderivative:

$$\ker(D_-) = \left\{ \bar{\Phi}, \bar{\Xi}, \Lambda, \Phi_L, (D_+ \bar{\Theta}), (\bar{D}_+ \bar{\Pi}), \left\{ \bar{D}_+^{(D_+ \text{preceding})} \right\}, D_- (\text{anything}) \right\}; \quad (2.30a)$$

$$\ker(D_+) = \left\{ \bar{\Phi}, \Xi, \Upsilon, \Phi_R, (D_- \bar{\Theta}), (\bar{D}_- \Pi), \left\{ \bar{D}_-^{(D_- \text{preceding})} \right\}, D_+ (\text{anything}) \right\}; \quad (2.30b)$$

$$\ker(\bar{D}_+) = \left\{ \Phi, \bar{\Xi}, \Upsilon, \bar{\Phi}_R, (\bar{D}_- \Theta), (D_- \bar{\Pi}), \left\{ \bar{D}_-^{(D_- \text{preceding})} \right\}, \bar{D}_+ (\text{anything}) \right\}; \quad (2.30c)$$

$$\ker(\bar{D}_-) = \left\{ \Phi, \Xi, \Lambda, \bar{\Phi}_L, (\bar{D}_+ \Theta), (D_+ \Pi), \left\{ \bar{D}_+^{(D_+ \text{preceding})} \right\}, \bar{D}_- (\text{anything}) \right\}. \quad (2.30d)$$

2.6. Other (super)constraints

Having listed the simple first-order and second-order superconstraints (2.1) and (2.19), respectively, it should be clear that even this is *by far* not all there is. One may also define

constrained superfields by means of less simple superderivative equations. Let us denote the sixteen components of a so far unconstrained superfield:

$$\Omega = \begin{pmatrix} & & W_{=} \\ & \omega_{-} & W & o_{-} \\ w & \varpi_{-} & W_{\mp} & \tilde{o}_{-} & \mathcal{O} \\ & \omega_{+} & W_{\pm} & o_{+} \\ & \varpi_{+} & W & \tilde{o}_{+} \\ & & W_{\mp} & \end{pmatrix} . \quad (2.31)$$

Yet higher order superconstraints

Having examined first and second order (simple) superconstraints in some detail, the inquisitive Reader will naturally wonder about third order superconstraints. The second order constraints examined in § 2.5 seem to produce merely a technically more involved cousins of the ‘minimal’ haploid superfields 2.1. Whereas the ‘minimal’ superfields are constrained by *two* simple, first order superconstraints which set three quarters of the component fields to zero, the non-minimal ones are constrained by a single, second-order superconstraint which sets only one quarter of the component fields to zero; the balance of a half of the component fields (eight of them) present in a non-minimal superfield but not in a ‘minimal’ one are the ‘extra’ component fields which in known applications turn out to be non-physical⁶⁾.

Naïvely then, we expect a cubic (simple) superconstraint to set only an eighth (two) of the component fields to zero. Also, we expect several of the non-vanishing components turning out to be auxiliary (with algebraic, *i.e.*, non-differential equations of motion), the precise identification possibly strongly depending on the choice of the Lagrangian density.

This turns out to be only partially true; in particular, the first expectation turns out to be wrong.

Let us examine the case

$$[D_{-}, \bar{D}_{-}]D_{+}\Omega = 0 . \quad (2.32)$$

Easily, by projecting on the lowest component of (2.32), this includes the simple statement that

$$0 \stackrel{!}{=} [D_{-}, \bar{D}_{-}]D_{+}\Omega \stackrel{\text{def}}{=} 4\sqrt{2}o_{-} , \quad (2.33)$$

so $o_{-} = 0$. Projecting with D_{-} , $\bar{D}_{-}$ and $\bar{D}_{-}$ (D_{+} of course trivially annihilates the whole superconstraint), we obtain the next three component constraints:

$$0 \stackrel{!}{=} -4i(\partial_{=}W) , \quad 0 \stackrel{!}{=} -4(\partial_{=}W_{\pm}) , \quad 0 \stackrel{!}{=} 4i(\partial_{\mp}W_{=}) - 8\mathcal{O} . \quad (2.34)$$

These imply, respectively, that W and $W_{\pm}$ are left-movers, and that the ‘top’ component field, $\mathcal{O}$, is not independent, but equals $\mathcal{O} = \frac{i}{2}(\partial_{\mp}W_{=})$. Continuing this, with the quadratic and cubic superdifferential operators from Fig. 1 which do not include D_{+} , we obtain four

⁶⁾ Recall that the property of a component field being physical, *i.e.*, propagating (with a differential equation of motion) depends most crucially on the choice of the Lagrangian density!

more constraints. The net effect of the superconstraint (2.32) is then summarized by writing (in a hopefully obvious doublet notation):

$$\partial_{=}^2 \{\omega_+, W_{\mp}\} = \{0, \frac{i}{2}(\partial_{=}^2 \partial_{\mp} w)\} , \quad \{w, \varpi_+\} \text{ is free}, \quad (2.35a)$$

$$\partial_{=} \{W_{\pm}, \tilde{o}_+\} = \{0, \frac{i}{2}(\partial_{=} \partial_{\mp} \varpi_-)\} , \quad \{\varpi_-, W\} \text{ is free}, \quad (2.35b)$$

$$\partial_{=} \{W, o_+\} = \{0, \frac{i}{2}(\partial_{=} \partial_{\mp} \omega_-)\} , \quad \{\omega_-, W_{\mp}\} \text{ is free}, \quad (2.35c)$$

$$\{o_-, \mathcal{O}\} = \{0, \frac{i}{2}(\partial_{\mp} W_{=})\} , \quad \{W_{=}, \tilde{o}_-\} \text{ is free}. \quad (2.35d)$$

Thus, half of the component fields, in the right column of Eqs. (2.35), are untouched by the superconstraint (2.32); a quarter of the component fields obey a differential relationship to a half of the free component fields. Finally, an eighth of the component fields ($W_{\pm}$ and W) become left-movers, one component field (ω_+) an ‘almost left-mover’, and only one component field, o_- outright vanishes. The fact that these results can be organized into the doublet formation of (2.35) is a simple consequence that these doublets form $\bar{\varsigma}^+$ -dependent $(0,1)$ -superfields, $\bar{D}_+$ being the one superderivative not appearing in Eq. (2.32).

The straightforward expectations stated above did not turn out to be true simply because the superderivative operators do not all (anti)commute, whereupon the counting of independent degrees of freedom remaining unconstrained by a superconstraint is far from (that) simple. Obviously, there are four (essentially distinct) cubic simple superconstraints, and a single quartic one.

Having illustrated this point and with the utility of these superfields entirely unexplored, we now turn to some other, more radically novel possibilities.

Non-simple superconstraints

The (again) first-order superconstraint

$$(D_- - \bar{D}_-)\Omega = 0 \quad (2.36)$$

defines an independent pair of haploid (in fact, lefton) superfields. To see this, switch to the $(0,2)$ -superfield notation as in Eq. (2.11) and (2.24), and define

$$\mathring{w} \stackrel{\text{def}}{=} \Omega \rfloor , \quad \mathring{\omega}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} D_- \Omega \rfloor , \quad \mathring{\varpi}_- \stackrel{\text{def}}{=} \frac{1}{\sqrt{2}} \bar{D}_- \Omega \rfloor , \quad \mathring{W}_- \stackrel{\text{def}}{=} \frac{1}{4} [D_-, \bar{D}_-] \Omega \rfloor , \quad (2.37)$$

where $\rfloor$ denotes setting $\varsigma^- = 0 = \bar{\varsigma}^-$ but leaving $\varsigma^+, \bar{\varsigma}^+$ intact. Taking the same projections of the superconstraint (2.36), we obtain the $(0,2)$ -superconstraints, which almost immediately yield:

$$\mathring{\varpi}_- = \mathring{\omega}_- , \quad \mathring{W}_- = 0 , \quad (\partial_{=} \mathring{w}) = 0 , \quad \text{and} \quad (\partial_{=} \mathring{\omega}_-) = 0 . \quad (2.38)$$

Thus, the superfield obeying the superconstraint (2.36) consists of a spin 0 left-moving (haploid) superfield, $\mathring{w} = (w; \omega_+, \varpi_+; W_{\mp})$, and two identical copies of a spin $+\frac{1}{2}$ left-moving (haploid) superfield, $\mathring{\omega}_- = (\omega_-; W, W_{\mp}; o_+)$.

Note that the complex conjugate of Eq. (2.36) merely states that $\bar{\Omega}$, the hermitian conjugate of Ω , satisfies the identical superconstraint, not that Ω is real.

Multi-field constraints

Of course, there is no *a priori* reason to restrict oneself to (super)constraints involving only one superfield. For example, given a chiral superfield, Φ , and a twisted-chiral one, Ξ , the superconstraint

$$D_+ \Phi - \bar{D}_+ \Xi \stackrel{!}{=} 0 \quad (2.39)$$

clearly identifies the fermionic component fields $\psi_+ = \chi_+$ (by straightforward projection). On projecting with D_- :

$$D_-(D_+ \Phi - \bar{D}_+ \Xi) = 2(F - X) \stackrel{!}{=} 0 , \quad (2.40)$$

we find that the ‘auxiliary’ fields F and X are also identified. Projecting further on the remaining six components (using all the superderivative operators from Fig. 1 which do not include trivially acting $\bar{D}_-$), we find that the superconstraint (2.39) includes the above identifications, but also forces all of the components of both Φ and Ξ to be right-moving! That is, Eq. (2.39) *includes* also the constraints $(\partial_+ \Phi) = 0 = (\partial_+ \Xi)$.

What naïvely started out as identifying only some two subsets of two superfields, turned out to have a quite more restrictive end. It is thus imperative to always expand all the superconstraint(s) in their component constraints, so as to identify their full effect.

Inhomogeneous superconstraints

Next, one may wish to attempt mixed-order (inhomogeneous in the D ’s) superconstraints. So, consider

$$(\mathbb{1} - \tfrac{1}{2} D_- D_+) \Omega = 0 , \quad (2.41)$$

which, a bit surprisingly, turns out to annihilate the whole superfield! To see this, we switch to an ‘antichiral’ (1,1)-superfield notation, expanding in the ς ’s, but keeping the component (1,1)-superfields still dependent on the $\bar{\varsigma}$ ’s. Define

$$\mathring{w} \stackrel{\text{def}}{=} \Omega \rfloor , \quad \mathring{\omega}_\pm \stackrel{\text{def}}{=} \tfrac{1}{\sqrt{2}} D_\pm \Omega \rfloor , \quad \mathring{W} \stackrel{\text{def}}{=} \tfrac{1}{2} D_- D_+ \Omega \rfloor , \quad (2.42)$$

where $\rfloor$ denotes setting $\varsigma^\pm = 0$ but leaving $\bar{\varsigma}^\pm$ intact. Projecting in the similar fashion the superconstraint (2.42), we obtain, respectively:

$$\mathring{w} = \mathring{W} , \quad \mathring{\omega}_\pm = 0 , \quad \text{and} \quad \mathring{\bar{W}} = 0 . \quad (2.43)$$

Combined, they set the whole superfield obeying (2.42) to zero.

Another easy (and far less trivial) example is

$$(\mathbb{1} - \tfrac{1}{4} [D_-, \bar{D}_-][D_+, \bar{D}_+]) \Omega = 0 , \quad (2.44)$$

which does not annihilate Ω , but restricts all the component fields to solutions of simple differential equations:

$$\begin{aligned} (1 - \partial_- \partial_+) \{ W_\mp, W_\pm, \mathcal{O} \} &= 0 , & (1 + \partial_- \partial_+) \{ W, \bar{W} \} &= 0 , \\ (1 - \partial_-^2 \partial_+^2) \{ w; \omega_\pm, \varpi_\pm; W_-, W_+; o_\pm, \tilde{o}_\pm \} &= 0 , \end{aligned} \quad (2.45)$$

where, of course, $(1 - \partial_-^2 \partial_+^2) \equiv (1 - \partial_- \partial_+)(1 + \partial_- \partial_+)$. In this case, all component fields are restricted ‘on the mass shell’, their mass being ± 1 .

Families of superconstraints

We next consider a notion that appears to be novel in supersymmetry, but has become fairly familiar in superstring theory: we construct families of constraints.

A small modification of the superconstraint (2.41):

$$(\mathbb{1} - \tfrac{1}{2}\alpha D_- D_+) \Omega = 0, \quad \alpha = \text{const.}, \alpha \neq 0, \infty, \quad (2.46)$$

now involves a parameter α . As it turns out, this superconstraint family still annihilates the superfield Ω and so introduces nothing new. The limiting value $\alpha = 0$ may thus in fact be included without any change in the enforced vanishing of Ω . However, the limiting case $\alpha = \infty$ turns Ω into a non-minimal antichiral superfield—far from vanishing.

On the other hand, consider the pair of superconstraints:

$$[\sin(\tfrac{1}{2}\beta) D_+ - \cos(\tfrac{1}{2}\beta) \bar{D}_+] \Omega = 0 = \bar{D}_- \Omega. \quad (2.47)$$

Clearly, the special cases $\beta = 0$ and $\beta = \pi$ constrain Ω to be a chiral or a twisted-chiral superfield, respectively. That is, the family of superconstraints (2.47), parametrized by the angle β , interpolates *continuously* between chiral and twisted-chiral superfields! This may seem perplexing, since the swap $C_+ : \Phi \leftrightarrow \Xi$ was at first regarded as a *discrete* transformation, whereas it is now shown to be a continuous $SO(2) \simeq U(1)$ transformation. In fact, this transformation may be identified with the $U(1)_L$ subgroup of the general $SU(2)_+$ group acting on the doublet of supercoordinates $\varsigma^+, \bar{\varsigma}^+$, which in turn is the ‘left-handed’ half of $SU(2)_+ \times SU(2)_-$, the maximal compact group of transformations of the supercoordinates $\varsigma, \bar{\varsigma}$ which commutes with the Lorentz symmetry; see § 1.3.

It is then natural to ask what does Ω of Eq. (2.46) become for angles $\beta \neq 0, \pi$. Straightforwardly, the superconstraint system (2.47) then defines a pair of chiral leftons (2.3a), (Φ_R, Φ_{R+}) . In the limiting cases, we find

$$\Phi = \left\{ \begin{array}{ccccc} (\phi, \psi_-) & \xleftarrow{\quad} & \Phi_R = (\phi_R, \psi_{R-}) & \xrightarrow{\quad} & (x, \xi_-) \\ & \xleftarrow{0 \leftarrow \beta} & & \xrightarrow{\beta \rightarrow \pi} & \\ (\psi_+, F) & \xleftarrow{\quad} & \Phi_{R+} = (\phi_{R+}, \psi_{R+-}) & \xrightarrow{\quad} & (\chi_+, X) \end{array} \right\} = \Xi. \quad (2.48)$$

The ‘upper halves’ (as displayed here) of Φ and Ξ are easy to map to each other, and in fact identify: both pairs of component fields are projected the same way. In contradistinction, the mapping of the ‘lower halves’, $(\psi_+, F) \leftarrow (\phi_{R+}, \psi_{R+-}) \rightarrow (\chi_+, X)$, is non-trivial and involves the twist $D_+ \leftrightarrow \bar{D}_+$. The relation (2.48) then shows how the pair of chiral rightons interpolates between a chiral and a twisted-chiral superfield.

Relation to dualities

Since the chiral $\leftrightarrow$ twisted-chiral mapping may be identified as the root of the ‘mirror map’ [14], the interpolation (2.48) may provide a more complete description. While a detailed study of this is beyond our present scope, we cannot help noticing that that is but one element of the discrete symmetry $2S_4$ identified in § 1.3. Another operation,

the left-handed (or the right-handed, of course) parity, denoted $\mathbf{p}$ in (1.10c, d), but when acting also on the world-sheet coordinates, was identified in Ref. [15] as the root of the Type IIA $\leftrightarrow$ Type IIB duality. With a total of 47 nontrivial operations, the group of discrete symmetries $2S_4$ from § 1.3 seems to be a pretty good candidate for seeking roots of other dualities in (super)string, M- and F-theories.

Fibration of superconstraints

Let us, however, return to the 1-parameter family of superconstraints (2.46). In retrospect, it is almost trivial, since all values of $\alpha \neq \infty$ lead to $\Omega = 0$. Exceptionally, at $\alpha = \infty$, $\Omega \neq 0$ and is in fact a non-minimal antichiral superfield! That is, the 1-dimensional parameter space $\{\alpha \neq \infty\} \simeq \mathbb{R}^1$ is fibered with identical fibres, parametrized by the identical constrained superfield, $\Omega = 0$. Compactifying $\{\alpha \neq \infty\} \simeq \mathbb{R}^1$ into $\{\alpha\} \simeq S^1$, by including the $\alpha = \infty$ point, the fibration becomes quite non-trivial. One can think of the total space of this fibration as a simple cover of S^1 (parametrized by the constant and trivial value $\Omega=0$), with one exceptional set, E , at $\alpha = \infty$, where E is parametrized by the non-minimal antichiral superfield satisfying $D_- D_+ \Omega = 0$.

A bit more interestingly perhaps, the 1-parameter family of superconstraints (2.47) is clearly non-trivial, since different values of α lead to different (constrained) superfields. Again, however, the parameter space, $\{\alpha\} \simeq S^1$ is fibred mostly by a pair of chiral rightons. At the two special points, $\alpha = 0, \pi$ however, this 2 + 2-dimensional fibre, (Φ_R, Φ_{R+}) , fuses into a single 4-dimensional one, Φ and Ξ , respectively. In fact, it is easy to turn (2.47) into a fibration over a (complex!) $\mathbb{P}^1$:

$$(z_1 D_+ - z_2 \bar{D}_+) \Omega = 0 = \bar{D}_- \Omega , \quad (2.49)$$

where the two complex numbers $(z_1, z_2) \simeq (\lambda z_1, \lambda z_2)$ parametrize a complex projective (parameter) space, $\mathbb{P}^1$. Note that the invariance under rescaling by a non-zero finite complex parameter, λ , arises as the obvious symmetry of the homogeneous equation $(z_1 D_+ - z_2 \bar{D}_+) \Omega = 0$. The chiral and twisted-chiral superfields, Φ, Ξ , are then seen as special fibres (the North and the South poles, respectively) in a non-trivial fibration of the righton doublet (Φ_R, Φ_{R+}) over $\mathbb{P}^1$.

Non-linear superconstraints

Whereas the simple modification (2.46) of the superconstraint (2.41) produced an (almost) trivial outcome, a further but still simple modification

$$(\mathbb{1} - \tfrac{1}{2} f(\Omega) D_- D_+) \Omega = 0 , \quad (2.50)$$

produces a rather more interesting situation. To see this, switch back to the (1,1)-superfield notation, expanding in the ς 's, but leaving the component fields still superfunctions of the $\bar{\varsigma}$'s. Using the definitions in (2.42) and writing $\mathring{f}^{(n)} \equiv f^{(n)}(\mathring{w})$, we find that the superconstraint (2.50) includes

$$\begin{aligned} \mathring{w} - \mathring{f} \mathring{W} &\stackrel{!}{=} 0 , & \mathring{\omega}_{\pm} [\mathring{f} - \mathring{w} \mathring{f}'] &\stackrel{!}{=} 0 , \\ \mathring{w} ([\mathring{f} - \mathring{w} \mathring{f}'] + \mathring{\omega}_- \mathring{\omega}_+ \mathring{f}'' \mathring{f}) &\stackrel{!}{=} 0 . \end{aligned} \quad (2.51)$$

The first constraint simply sets $\mathring{W} = \mathring{w}/\mathring{f}$. With the latter two, this then enforces $\Omega = (\mathring{w}; \mathring{\omega}_\pm; \mathring{w}/\mathring{f})$, where:

$$\mathring{w}, \mathring{\omega}_\pm = \begin{cases} \text{free} & \text{if } f(\Omega) = \Omega/W_0, \\ 0 & \text{if } f(\Omega) \neq \Omega/W_0, \end{cases} \quad W_0 \stackrel{\text{def}}{=} \lim_{\Omega \rightarrow 0} (\Omega/f(\Omega)) . \quad (2.52)$$

Again, we have a (degenerate) fibration: over the *parameter space of the function* f , we generically have the constant $\Omega = (0; 0, 0; W_0)$, while at the special subset where f becomes a constant multiple of its argument, we have the unconstrained but only $\frac{3}{4}$ - independent superfield, $\Omega = (\mathring{w}; \mathring{\omega}_\pm; \mathring{w}/\mathring{f})$ which, in the limit $f \rightarrow \infty$ becomes a non-minimal antichiral superfield, as found above. Recall that $\mathring{w}, \mathring{\omega}_\pm$ in turn are otherwise unconstrained $\bar{\varsigma}$ -dependent $(1, 1)$ -superfields.

Other non-homogeneous superderivative constraints may produce less trivial results and define other possibly interesting constrained superfields. Reasons for using such more complicated superconstraints are both far from obvious and model-dependent ⁷⁾. Whilst one is tempted to dismiss such more complicated constrained superfields on grounds of physical equivalence to (some combination of) ‘minimal’ haploid and quartoid superfields at the cost of greater technical complexity, it may well be that their *joint* use offers some distinct advantage, as does the use of ‘minimal’ *together* with non-minimal superfields [13]. More than a mere mention of this possibility, however, is beyond our present scope.

Discrete constraints

Finally, the imposition of reality as a constraint on a superfield is probably the best known type of constraint: $V^\dagger \stackrel{!}{=} V$ defines a real superfield, used for Yang-Mills type gauge vectors [6,7]. Note that the hermiticity of the entire superfield *does not* imply the hermiticity of the component fields. In fact, $V^\dagger \equiv \mathbf{C}(V) \stackrel{!}{=} V$ implies that, *e.g.*, the lowest component is real, but the $\frac{1}{2}D_-D_+V|$ and $\frac{1}{2}\bar{D}_+\bar{D}_-V|$ components are imaginary.

In view of our analysis of discrete symmetries of the 2-dimensional (2,2)-superspacetime, it is clear that there are many, many more such constraints possible. For example, $\mathbf{P}(W) \stackrel{!}{=} W$ defines a left-right (parity) symmetric, *i.e.*, scalar superfield; $\mathbf{C}_-(Y) \stackrel{!}{=} Y$ imposes a reality condition but only on the $\varsigma^-, \bar{\varsigma}^-$ -components; *etc.* There being 47 non-trivial elements of $2S_4$, there are 47 such types of constraints possible in the 2-dimensional (2,2)-superspacetime, and they can of course be combined with the superdifferential operators to produce some quite formidably looking constraints.

Again, an exhaustive analysis of all possibilities is clearly out of hand, and presently we remain content with the ‘minimal’ haploid superfields, and a brief involvement of their non-minimal counterparts.

⁷⁾ I thank Joe Polchinski for discussions on this point, and sharing some early results of his work with Simeon Hellerman on some superconstraint equations generalizing the present Eq. (2.49) into a fully non-linear (albeit holomorphic) superconstraints in Ω . Their work focuses on applications in supersymmetric quantum mechanics. The present models can be dimensionally reduced to such theories, albeit with some possible loss of generality, just as dimensional reduction of 4-dimensional models fails to capture most of the diversity presented herein.

3. Lagrangian Densities

Once the superfields have been defined, we can turn to their dynamics, to be governed by a Lagrangian density. As it turns out, 2-dimensional (2,2)-supersymmetric models have an unprecedented rich set of choices for the Lagrangian density, and consequently for the dynamics.

3.1. Unconstrained formalism

Owing to the nilpotency of the superderivatives, the complex ambidexterous constrained superfields can be expressed as superderivatives of unconstrained ones [7]. For example, given an unconstrained complex superfield, Ω , the following are ‘minimal’ haploid superfields:

$$(\bar{D}_+ \bar{D}_- \Omega) \text{ is chiral,} \quad i.e., \text{ it obeys Eqs. (2.1a) ,} \quad (3.1a)$$

$$(D_+ D_- \Omega) \text{ is antichiral,} \quad i.e., \text{ it obeys Eqs. (2.1b) ,} \quad (3.1b)$$

$$(D_+ \bar{D}_- \Omega) \text{ is twisted-chiral,} \quad i.e., \text{ it obeys Eqs. (2.1c) ,} \quad (3.1c)$$

$$(\bar{D}_+ D_- \Omega) \text{ is twisted-antichiral,} \quad i.e., \text{ it obeys Eqs. (2.1d) .} \quad (3.1d)$$

Similarly, given *two* unconstrained complex superfields, $\Omega^\pm$, the following define non-minimal haploid superfields:

$$(\bar{D}_+ \Omega^+ + \bar{D}_- \Omega^-) \text{ is NM-chiral,} \quad i.e., \text{ it obeys Eqs. (2.19a) ,} \quad (3.2a)$$

$$(D_+ \Omega^+ + D_- \Omega^-) \text{ is NM-antichiral,} \quad i.e., \text{ it obeys Eqs. (2.19b) ,} \quad (3.2b)$$

$$(D_+ \Omega^+ + \bar{D}_- \Omega^-) \text{ is NM-twisted-chiral,} \quad i.e., \text{ it obeys Eqs. (2.19c) ,} \quad (3.2c)$$

$$(\bar{D}_+ \Omega^+ + D_- \Omega^-) \text{ is NM-twisted-antichiral,} \quad i.e., \text{ it obeys Eqs. (2.19d) .} \quad (3.2d)$$

Amusingly, both the ‘minimal’ and the non-minimal *unidexterous* superfields elude a similarly easy description in terms of unconstrained superfields.

This then defines an ‘unconstrained’ formalism where each of the constrained superfields (2.1a–d) and (2.19a–d) is meticulously replaced by its proxy (3.1) and (3.2), respectively. Quantization is now straightforward in terms of path-integrals over the components of the unconstrained superfields Ω used in (3.1), and $\Omega^\pm$ used in (3.2). This approach is however unavoidably beset with a substantially increased amount of sheer algebra, and will not be pursued herein.

3.2. Constrained formalism

With a collection of several constrained superfields of the types defined in Eqs. (2.5), we now seek the most general Lagrangian density (subject to some hopefully plausible restrictions). Consider an analytic function, f , of some of the haploid superfields (2.1). In general [6,7], the Berezin integral of this f over only those $\varsigma, \bar{\varsigma}$ ’s on which f does depend will transform under a general supersymmetry transformation into a total derivative. A world-sheet integral of such a Berezin integral is then, up to world-sheet boundary terms,

invariant under supersymmetry. When considering only world-sheets without a boundary, as we do, such Berezin integrals provide supersymmetric Lagrangian densities.

With the benefit of hindsight, we discuss the details of the various Lagrangian density terms starting with those which require the most number of Berezin integrations. Also, for now, the quantization issues pertaining to the unidexterous (super)fields will be completely ignored, and all expressions will use fully constrained (super)fields.

Finally, while any superfield can be assigned overall spin as discussed in §2.4, for reasons that will soon become obvious, we consider endowing only a subset of the unidexterous superfields with overall spin. We write $\Lambda = (\Lambda^{\hat{a}}, \Lambda^{\hat{a}\pm})$, where the superfields counted by the index $\hat{a}$ remain as defined in (2.1e), while those indexed by $\tilde{a}$ satisfy $j_3(\Lambda^{\pm}) = 1$, so that $j_3\{\ell^{\pm}, \lambda_{+}^{\pm}, \tilde{\lambda}_{+}^{\pm}, L_{\pm}^{\pm}\} = \{1, \frac{1}{2}, \frac{1}{2}, 0\}$. A subset of the rightons is similarly endowed with overall spin -1 : $\Upsilon = (\Upsilon^{\hat{i}}, \Upsilon^{\hat{i}\pm})$.

The general Lagrangian density

The most general manifestly supersymmetric Lagrangian density is of the following form:

$$L = \frac{1}{2} \int d\zeta^{-} \Psi^{+} + h.c. \quad + \quad \frac{1}{2} \int d\zeta^{+} \Psi^{-} + h.c. \quad (3.3a)$$

$$+ \frac{1}{2} \int d^2\zeta \, W + h.c. \quad + \quad \frac{1}{2} \int d^2\zeta' \, \Sigma + h.c. \quad (3.3b, c)$$

$$+ \frac{1}{2} \int d^2\zeta^{-} N^{\pm} + h.c. \quad + \quad \frac{1}{2} \int d^2\zeta^{\pm} \tilde{N}^{\pm} + h.c. \quad (3.3d, e)$$

$$+ \frac{1}{4} \int d^2\zeta^{-} d\zeta^{+} M^{+} + h.c. \quad + \quad \frac{1}{4} \int d^2\zeta^{\pm} d\zeta^{-} \tilde{M}^{-} + h.c. \quad (3.3f, g)$$

$$+ \frac{1}{4} \int d^4\zeta \, K + h.c. \quad . \quad (3.3h)$$

Recall that fermionic integration is equivalent to (covariant) superderivative(s), followed by the projection on the ‘body’, $\zeta^{\pm}=0=\bar{\zeta}^{\pm}$:

$$L = \frac{1}{2} [D_{+}\Psi^{+}]| + h.c. \quad + \quad \frac{1}{2} [D_{-}\Psi^{-}]| + h.c. \quad (3.4a)$$

$$+ \frac{1}{2} [D_{-}D_{+}W]| + h.c. \quad + \quad \frac{1}{2} [D_{-}\bar{D}_{+}\Sigma]| + h.c. \quad (3.4b, c)$$

$$+ \frac{1}{4} [[D_{+}, \bar{D}_{+}]N^{\pm}]| + h.c. \quad + \quad \frac{1}{4} [[D_{-}, \bar{D}_{-}]\tilde{N}^{\pm}]| + h.c. \quad (3.4d, e)$$

$$+ \frac{1}{8} [[D_{+}, \bar{D}_{+}]D_{-}M^{+}]| + h.c. \quad + \quad \frac{1}{8} [[D_{-}, \bar{D}_{-}]D_{+}\tilde{M}^{-}]| + h.c. \quad (3.4f, g)$$

$$- \frac{1}{16} [[D_{-}, \bar{D}_{-}][D_{+}, \bar{D}_{+}]K]| \quad . \quad (3.4h)$$

For the above expressions to be supersymmetric, it is necessary and sufficient for each integrand to be annihilated by all covariant superderivatives except those which are equivalent to the integration operator. For example, Ψ^{+} should be a chiral lefton, annihilated by all superderivatives except D_{+} , which appears explicitly in (3.4a), replacing the integral $\int d\zeta^{-}$ in (3.3a). Similarly, W has to be chiral, Σ twisted-chiral, M^{+} has to be annihilated by $\bar{D}_{-}$, *etc.* K is always chosen to be real, so that no ‘+h.c.’ is necessary.

The $\int d\varsigma W + h.c.$ terms are frequently called the ‘F-terms’ as the $\int d\varsigma$ integral projects on the $\varsigma^+\varsigma^-$ -component in W , typically called F in a chiral superfield. We will extend the name to all other two-fermionic integrals in (3.3b, c), *i.e.*, (3.4b, c). The ς^4 component being typically called D in a general superfield, the $\int d^4\varsigma$ -term (3.3e), *i.e.*, (3.4e) is typically called the ‘D-term’. In a similar vein, we call the terms in (3.3a), *i.e.*, (3.4a) the ‘ ψ -terms’, and those in (3.3d), *i.e.*, (3.4d) the ‘ λ -terms’.

3.3. D-term

We begin with the most generally $\varsigma^\pm, \bar{\varsigma}^\pm$ -dependent set of terms, those that depend on all four of them, and so appear under the $\int d^4\varsigma$ integral (1.7). Since non-minimal superfields (2.19) do depend on all four fermionic coordinates, this is in fact the only type of terms in which they can appear (without a superderivative acting on them). The hasty Reader should be reminded that the *combined* use of ‘minimal’ and non-minimal haploid superfields does have its distinct advantages and interest (see, *e.g.*, Ref. [13]), whence here we include quartoid and both the ‘minimal’ and the (ambidexterous) non-minimal haploid superfields. The Reader should then have no problem extending the presented results by including the unidexterous non-minimal superfields, and also the ‘other’ superfields discussed in § 2.6, or indeed yet other ones, of their own design.

The D-term involving the haploid superfields is

$$L_K = \frac{1}{4} \int d^4\varsigma K(\Phi, \bar{\Phi}, \Xi, \bar{\Xi}, \Lambda, \Upsilon, \Theta, \bar{\Theta}, \Pi, \bar{\Pi}, \dots) , \quad (3.5)$$

where the ellipses denote non-minimal superfields, and $K(\dots)$ is required to be a real function for the Lagrangian (term) L_K to be hermitian. Note that this implies, *e.g.*, that $\overline{K_{,\mu\nu}} \stackrel{\text{def}}{=} \overline{\partial_\mu \partial_\nu K} = K_{,\nu\bar{\mu}}$ and $\overline{K_{,\alpha\bar{\beta}}} = K_{,\beta\bar{\alpha}}$ are Hermitian, but $K_{,ab}, K_{,ij}$ are merely symmetric, and all the mixed matrices are general. We do not, however, allow K to depend on superderivative superfields, as that would lead to higher derivative terms in the Lagrangian density.

Finally, as $\Lambda^\pm$ and $\Upsilon^\pm$ carry overall spin, and since K has spin-0 and it is not allowed to depend on superderivatives, it must be that K depends on the $\Lambda^\pm, \Upsilon^\pm$ only through their spinless products, $(\Lambda^\pm \Upsilon^\pm)$. Such products are characterized by a list of matrices, $\kappa_{\check{a}\check{i}}^m$, where m indexes all the linearly independent bilinear products, $B^m \stackrel{\text{def}}{=} (\Lambda^{\check{a}\pm} \kappa_{\check{a}\check{i}}^m \Upsilon^{\check{i}\pm})$. The matrices $\kappa_{\check{a}\check{i}}^m$ may be chosen so as to have all entries 0, except for the m^{th} one (counting say elements in row after row) which then equals 1. Then,

$$\frac{\partial}{\partial \Lambda^{\check{a}\pm}} = \kappa_{\check{a}\check{i}}^m \Upsilon^{\check{i}\pm} \frac{\partial}{\partial B^m} , \quad \text{and} \quad \frac{\partial}{\partial \Upsilon^{\check{i}\pm}} = \Lambda^{\check{a}\pm} \kappa_{\check{a}\check{i}}^m \frac{\partial}{\partial B^m} , \quad (3.6)$$

so that

$$\frac{\partial K_{,\dots}}{\partial \Lambda^{\check{a}\pm}} = \kappa_{\check{a}\check{i}}^m \Upsilon^{\check{i}\pm} K_{,m\dots} , \quad \text{and} \quad \frac{\partial K_{,\dots}}{\partial \Upsilon^{\check{i}\pm}} = \Lambda^{\check{a}\pm} \kappa_{\check{a}\check{i}}^m K_{,m\dots} . \quad (3.7)$$

With these, the Lagrangian density is straightforward to expand into component fields; see Appendix B.

3.4. λ -terms

We now seek *fermionic* functions which are annihilated by one of the four $D, \bar{D}$'s. Since Φ, Ξ , their $D_{\pm}$ -superderivatives, $\Lambda, (D_+\Lambda), (\bar{D}_+\Lambda)$ and $(\bar{D}_-\text{anything})$ are all annihilated by $\bar{D}_-$, the Berezin integral over $\varsigma^{\pm}, \bar{\varsigma}^{\pm}$ of any analytic function of these superfields is a suitable Lagrangian density term:

$$L_{M^+} = \frac{1}{4} \int d^2\varsigma d\bar{\varsigma}^- M^+(\Phi_L, \Phi_R, \bar{\Phi}_L; \Phi, \Xi, \Lambda; \dots) + h.c. , \quad (3.8)$$

as supersymmetry transforms it into a total derivative⁸⁾. Recall that the component fields (2.15) of the quartoid superfields $\Phi_L, \Phi_R, \bar{\Phi}_L, \bar{\Phi}_R$ are obtained from those (2.5a, b) of the chiral and antichiral haploid superfields. We can therefore lump Φ_L, Φ_R into Φ and $\bar{\Phi}_L, \bar{\Phi}_R$ into $\bar{\Phi}$ without any loss of generality; the interested Reader should have no difficulty re-introducing them by splitting the ranges: $\Phi \rightarrow (\Phi, \Phi_L, \Phi_R)$ and $\bar{\Phi} \rightarrow (\bar{\Phi}, \bar{\Phi}_L, \bar{\Phi}_R)$, and remembering that the quartoid superfields have only half the component fields of a haploid one. Here, the quartoid superfields are explicitly written only if they do not appear in pairs that comprise a haploid superfield.

Now, prohibiting as before higher derivative (negative dimensional coefficient) terms, and ensuring that M^+ has spin $+\frac{1}{2}$, it must be of the form:

$$M^+ = (D_- \Phi^\mu) M_\mu^L + (D_- \Xi^\alpha) M_\alpha^L + (D_+ \Lambda^{\check{a}\pm}) M_{\check{a}}^1 + (\bar{D}_+ \Lambda^{\check{a}\pm}) M_{\check{a}}^2 \quad (3.9a)$$

$$+ (\bar{D}_- \Phi^{\bar{\nu}}) M_{\bar{\nu}} + (\bar{D}_- \Xi^{\bar{\beta}}) M_{\bar{\beta}} + (\bar{D}_- \Upsilon^{\hat{i}}) M_{\hat{i}} + (\bar{D}_+ \Phi_L^{\check{\mu}}) \Lambda^{\check{a}\pm} M_{\check{a}\check{\mu}} \quad (3.9b)$$

$$+ (\bar{D}_- \Theta^M) M_M + (\bar{D}_- \Theta^{\bar{N}}) M_{\bar{N}} + (\bar{D}_- \Pi^A) M_A + (\bar{D}_- \Pi^{\bar{B}}) M_{\bar{B}} , \quad (3.9c)$$

$$+ (D_+ \Phi^\mu) \Lambda^{\check{a}\pm} M_{\check{a}\mu} + (D_+ \Lambda^{\hat{a}}) \Lambda^{\check{b}\pm} M_{\hat{a}\check{b}}^1 + (D_+ \Pi^A) \Lambda^{\check{b}\pm} M_{A\check{b}} \quad (3.9d)$$

$$+ (\bar{D}_+ \Xi^\alpha) \Lambda^{\check{b}\pm} M_{\alpha\check{b}} + (\bar{D}_+ \Lambda^{\hat{a}}) \Lambda^{\check{b}\pm} M_{\hat{a}\check{b}}^2 + (\bar{D}_+ \Theta^M) \Lambda^{\check{a}\pm} M_{\check{a}M} , \quad (3.9e)$$

where the M 's are all arbitrary analytic functions of the $\bar{\Phi}_L, \Phi, \Xi, \Lambda$, except for M_α^L, M_μ^L , which are depend only on $\bar{\Phi}_L, \Lambda$. These two terms are supersymmetric since, although they are not annihilated by $\bar{D}_-$ as the other are, they turn into total derivatives:

$$\bar{D}_- [(D_- \Phi^\mu) M_\mu^L] = (2i\partial_- \Phi^\mu) M_\mu^L = 2i\partial_- (\Phi^\mu M_\mu^L) , \quad (3.10)$$

which suffices.

The quick Reader may notice that the $(\bar{D}_- \dots)M$ -type terms in (3.9b, c) may be rewritten as D-terms (since $(\bar{D}_- \Phi^{\bar{\nu}}) M_{\bar{\nu}} = \bar{D}_- (\Phi^{\bar{\nu}} M_{\bar{\nu}})$, *etc.*), and could argue for their discarding here on the grounds that they are effectively included in (3.5). This would however induce a serious omission. Rather than advance a general and possibly obtuse

⁸⁾ This general argument rests on the identities (up to total deivatives) $Q_{\pm} \simeq D_{\pm}$ and $\bar{Q}_{\pm} \simeq \bar{D}_{\pm}$. Supersymmetry transformation is then calculated, up to total derivatives, by acting with the operator $\delta_{\epsilon} \simeq \epsilon \cdot D + \bar{\epsilon} \cdot \bar{D}$.

argument, consider the following simple example, with the first term in (3.9b) where $M_{\bar{\nu}} = \Phi^\mu M_{\mu\bar{\nu}}(\Phi)$:

$$\begin{aligned} & \frac{1}{8} \{ [D_+, \bar{D}_+] D_- (M_{\bar{\nu}}(\Phi)(\bar{D}_- \Phi^{\bar{\nu}})) + [\bar{D}_+, D_+] D_- ((D_- \Phi^\mu) \bar{M}_{\bar{\nu}\mu} \Phi^{\bar{\nu}}) \} | \\ & = \frac{1}{2} D^4 (\Phi^\mu M_{\mu\bar{\nu}}^H \Phi^{\bar{\nu}}) | - \frac{1}{4} \partial_=([D_+, \bar{D}_+] (\Phi^\mu M_{\mu\bar{\nu}}^A \Phi^{\bar{\nu}})) | , \end{aligned} \quad (3.11)$$

where $\bar{M}_{\bar{\nu}\mu} \stackrel{\text{def}}{=} (M_{\nu\bar{\mu}})^\dagger$ and

$$M_{\mu\bar{\nu}}^H \stackrel{\text{def}}{=} \frac{1}{2} (M_{\mu\bar{\nu}} + \bar{M}_{\bar{\nu}\mu}) , \quad M_{\mu\bar{\nu}}^A \stackrel{\text{def}}{=} \frac{1}{2i} (M_{\mu\bar{\nu}} - \bar{M}_{\bar{\nu}\mu}) \quad (3.12)$$

are hermitian and anti-hermitian, respectively. The first term in (3.11) contains, upon projection, the familiar ‘kinetic’ terms

$$\frac{1}{2} [(\partial_+ \phi^\mu)(\partial_- \phi^{\bar{\nu}}) + (\partial_- \phi^\mu)(\partial_+ \phi^{\bar{\nu}})] M_{\mu\bar{\nu}}^H , \quad (3.13)$$

whereas the second term in (3.11) includes the ‘torsion’ term:

$$\frac{1}{2} [(\partial_+ \phi^\mu)(\partial_- \phi^{\bar{\nu}}) - (\partial_- \phi^\mu)(\partial_+ \phi^{\bar{\nu}})] M_{\mu\bar{\nu}}^A . \quad (3.14)$$

The antihermitian matrix $M_{\mu\bar{\nu}}^A$ appearing here need not necessarily be associated with the ‘metric torsion’, which would be a derivative of $M_{\mu\bar{\nu}}^A$. In fact, $M_{\mu\bar{\nu}}^A$ may well be constant. However, it does produce an important term upon world-sheet integration:

$$\int d^2\sigma \frac{1}{2} [(\partial_+ \phi^\mu)(\partial_- \phi^{\bar{\nu}}) - (\partial_- \phi^\mu)(\partial_+ \phi^{\bar{\nu}})] M_{\mu\bar{\nu}}^A = \int d\phi^\mu \wedge d\phi^{\bar{\nu}} M_{\mu\bar{\nu}}^A \equiv \int \phi^* M_{[2]}^A , \quad (3.15)$$

the integral of the (the pull-back by the map ϕ of the) 2-form $M_{[2]}^A \equiv M_{\mu\bar{\nu}}^A d\phi^\mu \wedge d\phi^{\bar{\nu}}$ over the (image) of the world-sheet. This type of terms relates to *global* information about the field space and is essential in many superstring applications of 2-dimensional (supersymmetric) field theories.

Since total derivatives are routinely dropped, this important term would normally be lost. Thus, instead of rewriting the $\bar{D}_-(\dots)$ -terms in (3.9) as D-terms and then meticulously salvaging such and similarly interesting terms from the total derivative debris *en route*, it would seem preferable to include them manifestly in the Lagrangian density via the terms such as those in (3.9).

Notice also that the ‘metric’ $M_{\mu\bar{\nu}}^H$ defined in the first one of Eqs. (3.12) stems from a simple *choice* of $M_{\bar{\nu}}(\bar{\Phi}_L, \Phi, \Xi, \Lambda)$. In the general case, the ‘metric’ in the ‘kinetic’ term (3.13) is then a hermitian and analytic but otherwise unrestricted matrix function of the bosonic fields $\bar{\phi}_L, \phi, x, \ell$ and their conjugates.

Thus, the $(\bar{D}_- \dots)M$ -type terms in (3.9b, c) play a dual rôle: (1) they modify the D-terms, and (2) produce ‘topological’ terms such as (3.15). The remaining eleven terms in (3.9a–e) however cannot be reduced to D-terms and/or total derivatives; they produce totally new terms for the Lagrangian density.

By the same token, the other batch of λ -terms is obtained from

$$\widetilde{M}^- = (D_+ \Phi^\mu) \widetilde{M}_\mu^R + (D_+ \Xi^{\bar{\beta}}) \widetilde{M}_{\bar{\beta}}^R + (D_- \Upsilon^{i=}) \widetilde{M}_i^1 + (\bar{D}_- \Upsilon^{i=}) \widetilde{M}_i^2 \quad (3.9f)$$

$$+ (\bar{D}_+ \Phi^{\bar{\nu}}) \widetilde{M}_{\bar{\nu}} + (\bar{D}_+ \Xi^\alpha) \widetilde{M}_\alpha + (\bar{D}_+ \Lambda^{\hat{a}}) \widetilde{M}_{\hat{a}} + (\bar{D}_- \Phi_R^{\bar{\mu}}) \Upsilon^{i=} \widetilde{M}_{i\bar{\mu}} \quad (3.9g)$$

$$+ (\bar{D}_+ \Theta^M) \widetilde{M}_M + (\bar{D}_+ \Theta^{\bar{N}}) \widetilde{M}_{\bar{N}} + (\bar{D}_+ \Pi^A) \widetilde{M}_A + (\bar{D}_+ \Pi^{\bar{B}}) \widetilde{M}_{\bar{B}} , \quad (3.9h)$$

$$+ (D_- \Phi^\mu) \Upsilon^{i=} \widetilde{M}_{i\mu} + (D_- \Upsilon^i) \Upsilon^{j=} \widetilde{M}_{ij}^1 + (D_- \Pi^{\bar{B}}) \Upsilon^{j=} \widetilde{M}_{\bar{B}j} \quad (3.9i)$$

$$+ (\bar{D}_- \Xi^{\bar{\beta}}) \Upsilon^{j=} \widetilde{M}_{\bar{\beta}j} + (\bar{D}_- \Upsilon^i) \Upsilon^{j=} \widetilde{M}_{ij}^2 + (\bar{D}_- \Theta^M) \Upsilon^{i=} \widetilde{M}_{iM} , \quad (3.9j)$$

where now the $\widetilde{M}$.'s are all arbitrary analytic functions of the $\bar{\Phi}_R, \Phi, \Xi, \Upsilon$, except for $\widetilde{M}_\mu^R$ and $\widetilde{M}_{\bar{\beta}}^R$ which only depend on $\bar{\Phi}_R, \Upsilon$.

With these two functions (3.9),

$$L_M = \frac{1}{8} [[D_+, \bar{D}_+] D_- M^+] | + h.c. \quad (3.16a)$$

$$+ \frac{1}{8} [[D_-, \bar{D}_-] D_+ \widetilde{M}^-] | + h.c. \quad (3.16b)$$

is the most general set of λ -terms for the Lagrangian density; see also Appendix B.

3.5. *F-terms*

‘Superpotential terms’ (3.3b) are a familiar type of candidates for the Lagrangian density. For example, any analytic function of only the chiral superfields, Φ ’s, is itself a chiral superfield: $\bar{D}_\pm W(\Phi)=0$.

Ambidexterous superpotentials

Since all the superfields in (2.29a) satisfy the chiral pair of superconstraints (2.1a), any analytic function of the arguments listed in (2.29a),

$$W(\Phi, \Phi_L, \Phi_R; (\bar{D}_+ \Xi), \dots) ,$$

is than also chiral. Its hermitian conjugate, $\bar{W}$, is antichiral. Similarly,

$$\Sigma(\Xi, \Phi_R, \bar{\Phi}_L; (D_+ \Phi), \dots) ,$$

is than also twisted chiral, and its hermitian conjugate, $\bar{\Sigma}$, is twisted antichiral. Again, we lump Φ_L, Φ_R into Φ and $\bar{\Phi}_L, \bar{\Phi}_R$ into $\bar{\Phi}$, and write out explicitly the dependence on quartoid superfields only if this cannot be done, as in specifying the dependence of the coefficient functions in (3.9).

However, a few further restrictions are needed. For the putative Lagrangian density terms

$$\int d^2\zeta W + h.c. \quad \text{and} \quad \int d^2\zeta \Sigma + h.c. , \quad (3.17)$$

to be scalar densities, W and Σ have to be scalar densities themselves. Therefore, each term in their expansion over the superderivative superfields must have an even number of superderivatives. Finally, to prevent higher derivative and/or negative dimensional coefficient terms, we write W in the following form:

$$W = W_0(\Phi) + \frac{1}{2} W_{>}(\Phi; \dots) , \quad (3.18a)$$

and truncate $W_{>}$ to terms quadratic in superderivatives.

Now, among all the possible terms with a quadratic superderivative, *e.g.*, the term $\int d^2\zeta (\bar{D}_+\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}} + h.c.$ can be rewritten, without any loss of generality, as a corresponding λ -term (3.16):

$$\begin{aligned} \int d^2\zeta (\bar{D}_+\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}}(\Phi) &\equiv D_-D_+(\bar{D}_+\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}}(\Phi) = D_-D_+\bar{D}_+(\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}}(\Phi) , \\ &\simeq \frac{1}{2}[D_+, \bar{D}_+]D_- [(\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}}(\Phi)] , \end{aligned} \quad (3.19)$$

where we have dropped the total derivative $i\partial_+D_-[(\bar{D}_-\Phi^{\bar{\nu}})W_{\bar{\nu}}]$. In fact, the corresponding λ -term in (3.9b), $\int d^2\zeta d\bar{\zeta}^-(\bar{D}_-\Phi^{\bar{\nu}})M_{\bar{\nu}}$, is *more* general, as the coefficient function $M_{\bar{\nu}}$ may depend on $\bar{\Phi}_L, \Phi, \Xi, \Lambda$, whereas $W_{\bar{\nu}}$ depends only on Φ . A similar fate befalls many of the tentative F-terms, and only the following remain, not being reducible without loss of generality to any of the terms in (3.5) or (3.16):

$$\begin{aligned} W_{>} &= (\bar{D}_+\Lambda^{\check{a}\dagger})(\bar{D}_-\Upsilon^{\check{i}=})W_{\check{a}\check{i}} + (\bar{D}_+\Lambda^{\check{a}\dagger})(\bar{D}_+\Theta^M)W_{\check{a}M} \\ &\quad + (\bar{D}_-\Upsilon^{\check{i}=})(\bar{D}_-\Theta^M)W_{iM} + (\bar{D}_+\Theta^M)(\bar{D}_-\Theta^N)W_{MN} , \end{aligned} \quad (3.18b)$$

where the $W_{\cdot\cdot}$'s are arbitrary analytic functions of the Φ only. The canonical dimensions of $W_0(\Phi)$ and the coefficient functions in $W_{>}(\Phi)$ are all non-negative: $[W_0]=1$ (mass parameters) and $[W_{\cdot\cdot}]=0$ (dimensionless couplings). The omitted higher order superderivative terms would have had coefficient functions with negative dimensions. With the superpotential (3.18),

$$L_W = \frac{1}{2} \int d^2\zeta W(\Phi; \dots) + h.c. = \frac{1}{2}[D_-D_+W(\Phi; \dots)] + h.c. \quad (3.20)$$

is the most general superpotential Lagrangian density. In the absence of unidexterous superfields, L_W reduces to the well-known superpotential terms involving only the holomorphic function $W_0(\Phi)$ in (3.18).

Similarly,

$$L_{\Sigma} = \frac{1}{2} \int d^2\zeta \Sigma(\Xi; \dots) + h.c. = \frac{1}{2}[D_-\bar{D}_+\Sigma(\Xi; \dots)] + h.c. \quad (3.21)$$

is the most general twisted superpotential Lagrangian density, with the twisted superpotential being

$$\Sigma = \Sigma_0(\Phi) + \frac{1}{2}\Sigma_{>}(\Xi; \dots) , \quad (3.22a)$$

where

$$\begin{aligned} \Sigma_{>} &= (D_+\Lambda^{\check{a}\dagger})(\bar{D}_-\Upsilon^{\check{i}=})\Sigma_{\check{a}\check{i}} + (\bar{D}_+\Pi^A)(D_+\Lambda^{\check{b}\dagger})\Sigma_{A\check{b}} \\ &\quad + (D_-\Pi^A)(\bar{D}_-\Upsilon^{\check{i}=})\Sigma_{A\check{i}} + (D_+\Pi^A)(\bar{D}_-\Pi^B)\Sigma_{AB} , \end{aligned} \quad (3.22b)$$

where the $\Sigma_{\cdot\cdot}$'s are arbitrary analytic functions of the Ξ only. The canonical dimensions of $\Sigma_0(\Phi)$ and the coefficient functions in $\Sigma_{>}(\Phi)$ are all non-negative: $[\Sigma_0]=1$ (mass parameters) and $[\Sigma_{\cdot\cdot}]=0$ (dimensionless couplings).

Unidexterous superpotentials

By contrast, no (higher) superderivative of either of $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}$ satisfies the superconstraints (2.1e, f). Therefore, L_L (L_R), the lefton (righton) analogue of L_W and L_Σ , is constructed entirely from lefton (righton) superfields. Using the list (2.29e), we find that

$$\begin{aligned} L_L &= \frac{1}{2} \int d^2\varsigma^\pm N^\pm(\Lambda; \dots) + h.c. \\ &= \frac{1}{4} [[D_+, \bar{D}_+] N^\pm(\Lambda; \dots)] + h.c. \end{aligned} \quad (3.23)$$

is the lefton ‘superpotential’ term for the Lagrangian density. For this to be a scalar density, the function $N^\pm$ must transform as a vector of spin +1, and so must depend on the $\Lambda^\pm$ ’s (this is the reason for their introduction). Restricting again to terms in which the coefficient functions have non-negative dimensions, we find (omitting indices for brevity):

$$\begin{aligned} N^\pm &= \Lambda^\pm N. + \frac{1}{2} \Lambda^\pm (D_+ \bar{D}_+ \Lambda^\pm) N.^1 + \frac{1}{2} \Lambda^\pm (\bar{D}_+ D_+ \Lambda^\pm) N.^2 \\ &\quad + \frac{1}{2} (D_+ \Lambda^\pm) (D_+ \Lambda^\pm) N.^3 + \frac{1}{2} (D_+ \Lambda^\pm) (\bar{D}_+ \Lambda^\pm) N.^4 \\ &\quad + \frac{1}{2} (\bar{D}_+ \Lambda^\pm) (\bar{D}_+ \Lambda^\pm) N.^5 + \frac{1}{2} \Lambda^\pm \Lambda^\pm (D_+ \bar{D}_+ \Lambda) N.^1.. \\ &\quad + \frac{1}{2} \Lambda^\pm \Lambda^\pm (\bar{D}_+ D_+ \Lambda) N.^2.. + \frac{1}{2} \Lambda^\pm (D_+ \Lambda^\pm) (D_+ \Lambda) N.^3.. \\ &\quad + \frac{1}{2} \Lambda^\pm (D_+ \Lambda^\pm) (\bar{D}_+ \Lambda) N.^4.. + \frac{1}{2} \Lambda^\pm (\bar{D}_+ \Lambda^\pm) (D_+ \Lambda) N.^5.. \\ &\quad + \frac{1}{2} \Lambda^\pm (\bar{D}_+ \Lambda^\pm) (\bar{D}_+ \Lambda) N.^6.. + \frac{1}{2} \Lambda^\pm \Lambda^\pm (D_+ \Lambda) (D_+ \Lambda) N.^1... \\ &\quad + \frac{1}{2} \Lambda^\pm \Lambda^\pm (D_+ \Lambda) (\bar{D}_+ \Lambda) N.^2... + \frac{1}{2} \Lambda^\pm \Lambda^\pm (\bar{D}_+ \Lambda) (\bar{D}_+ \Lambda) N.^3... , \end{aligned} \quad (3.24a)$$

where the $N., \dots, N....$ ’s are all arbitrary functions of the Λ . Clearly, some of these have certain symmetries, implicit in the above expressions; *e.g.*, $N_{\bar{a}\bar{b}\bar{c}\bar{d}}^1 = +N_{\bar{b}\bar{a}\bar{c}\bar{d}}^1 = -N_{\bar{a}\bar{b}\bar{d}\bar{c}}^1$. That is, $N.^{1...} \rightarrow N.^{1}_{(\bar{a}\bar{b})[\bar{c}\bar{d}]}$. Also, the Reader may prefer to combine some of the terms, such as

$$\begin{aligned} &\frac{1}{2} \Lambda^\pm (D_+ \bar{D}_+ \Lambda^\pm) N.^1 + \frac{1}{2} \Lambda^\pm (\bar{D}_+ D_+ \Lambda^\pm) N.^2 \\ &= \frac{i}{4} \Lambda^\pm (\partial_\pm \Lambda^\pm) (N.^1 + N.^2) + \frac{1}{4} \Lambda^\pm ([D_+, \bar{D}_+] \Lambda^\pm) (N.^1 - N.^2) . \end{aligned} \quad (3.25)$$

Similarly for rightons,

$$\begin{aligned} L_R &= \frac{1}{2} \int d^2\varsigma^\mp \tilde{N}^\mp(\Upsilon; \dots) + h.c. \\ &= \frac{1}{4} [[D_-, \bar{D}_-] \tilde{N}^\mp(\Upsilon; \dots)] + h.c. \end{aligned} \quad (3.26)$$

where $N^\pm$ may depend on all of $\Upsilon, \Upsilon^\pm, (D_- \Upsilon), (\bar{D}_- \Upsilon)$ and $(D_- \bar{D}_- \Upsilon)$. Expanding this function in the superderivatives and the $\Upsilon^\pm$ ’s, and truncating the expansion so as to prohibit higher derivative terms and non-negative canonical dimension coefficients, we

find:

$$\begin{aligned}
\tilde{N}^= &= \Upsilon^= \tilde{N}^= + \frac{1}{2} \Upsilon^= (D_- \bar{D}_- \Upsilon^=) \tilde{N}^{1..} + \frac{1}{2} \Upsilon^= (\bar{D}_- D_- \Upsilon^=) \tilde{N}^{2..} \\
&+ \frac{1}{2} (D_- \Upsilon^=) (D_- \Upsilon^=) \tilde{N}^{3..} + \frac{1}{2} (D_- \Upsilon^=) (\bar{D}_- \Upsilon^=) \tilde{N}^{4..} \\
&+ \frac{1}{2} (\bar{D}_- \Upsilon^=) (\bar{D}_- \Upsilon^=) \tilde{N}^{5..} + \frac{1}{2} \Upsilon^= \Upsilon^= (D_- \bar{D}_- \Upsilon) \tilde{N}^{1..} \\
&+ \frac{1}{2} \Upsilon^= \Upsilon^= (\bar{D}_- D_- \Upsilon) \tilde{N}^{2..} + \frac{1}{2} \Upsilon^= (D_- \Upsilon^=) (D_- \Upsilon) \tilde{N}^{3..} \\
&+ \frac{1}{2} \Upsilon^= (D_- \Upsilon^=) (\bar{D}_- \Upsilon) \tilde{N}^{4..} + \frac{1}{2} \Upsilon^= (\bar{D}_- \Upsilon^=) (D_- \Upsilon) \tilde{N}^{5..} \\
&+ \frac{1}{2} \Upsilon^= (\bar{D}_- \Upsilon^=) (\bar{D}_- \Upsilon) \tilde{N}^{6..} + \frac{1}{2} \Upsilon^= \Upsilon^= (D_- \Upsilon) (D_- \Upsilon) \tilde{N}^{1..} \\
&+ \frac{1}{2} \Upsilon^= \Upsilon^= (D_- \Upsilon) (\bar{D}_- \Upsilon) \tilde{N}^{2..} + \frac{1}{2} \Upsilon^= \Upsilon^= (\bar{D}_- \Upsilon) (\bar{D}_- \Upsilon) \tilde{N}^{3..} ,
\end{aligned} \tag{3.24b}$$

where the $\tilde{N}^=, \dots, \tilde{N}^{6..}$'s are all arbitrary functions of the Υ .

3.6. ψ -terms

Finally, we come to the terms with the simplest $\varsigma, \bar{\varsigma}$ -dependence. As listed in (2.29g), $\Phi_L, (\bar{D}_+ \Lambda)$ and $(\bar{D}_+ D_+ \Lambda)$ satisfy the chiral lefton pair of superconstraints (2.3a), *i.e.*, they depend only on ς^- . Thus

$$L_{\Psi_L} = \int d\varsigma^- \Psi^+(\Phi_L; \bar{D}_+ \Lambda; \bar{D}_+ D_+ \Lambda) + h.c. \tag{3.27}$$

is a hermitian supersymmetric Lagrangian density term, given any analytic spin $+\frac{1}{2}$ function $\Psi^+(\Phi_L; \bar{D}_+ \Lambda; \bar{D}_+ D_+ \Lambda)$ and its hermitian conjugate⁹⁾.

The superfields $\Phi_L, \bar{D}_+ \Lambda$ and $\bar{D}_+ D_+ \Lambda$, as defined above (2.3) and (2.1), contain no positive spin component fields, which is why the $\Lambda^\pm$ have been introduced. For a Lagrangian density without higher derivative and negative dimensional coefficient functions, we truncate the expansion of Ψ^+ in $\bar{D}_+ \Lambda$ and $\bar{D}_+ D_+ \Lambda$, and obtain the list:

$$(\bar{D}_+ \Lambda^{\check{a}\pm}) \Psi_{\check{a}} , \quad \frac{1}{4} (\bar{D}_+ \Lambda^{\check{a}\pm}) (\bar{D}_+ D_+ \Lambda^{\check{b}\pm}) \Psi_{\check{a}\check{b}} , \quad + \frac{1}{4} (\bar{D}_+ \Lambda^{\check{a}\pm}) (\bar{D}_+ \Lambda^{\check{b}\pm}) (\bar{D}_+ \Lambda^{\check{c}}) \Psi_{\check{a}\check{b}\check{c}} , \tag{3.28}$$

where the $J_{\check{a}}, \Psi_{\check{a}\check{b}}$ and $\Psi_{\check{a}\check{b}\check{c}}$'s are arbitrary analytic functions of only the Φ_L . However, all of these turn out to be special cases of various terms from (3.24a). For example,

$$(\bar{D}_+ \Lambda^{\check{a}\pm}) \Psi_{\check{a}} \tag{3.29}$$

may be rewritten unambiguously (up to total derivatives) and without any loss of generality, as a unidexterous F-term:

$$\frac{1}{2} [D_+ (\bar{D}_+ \Lambda^{\check{a}\pm}) \Psi_{\check{a}}(\Phi_L)] + h.c. \cong \frac{1}{4} [[D_+, \bar{D}_+] \Lambda^{\check{a}\pm} \Psi_{\check{a}}(\Phi_L)] + h.c. \tag{3.30}$$

The same of course holds for its conjugate, and their parity mirrors, $\Psi^-(\Phi_R; \bar{D}_- \Upsilon; \bar{D}_- D_- \Upsilon)$ and its conjugate.

⁹⁾ Recall that the unidexterous superfields, Λ and Υ , while in general complex, *may be chosen* to be real or imaginary.

Note that it was necessary to introduce the superfields $\Lambda^\pm$ and $\Upsilon^\pm$ for there to be any unidexterous F-terms, (3.23) and (3.26). The Readers interested only in the ambidextrous haploid superfields (2.1a–d) need not concern themselves with (3.23) and (3.26), and the appropriate terms from the rest of (3.3).

3.7. Physical degrees of freedom

Note that *each term* in Eqs. (3.5), (3.16), (3.20), (3.21), (3.23) and (3.26) leads to a *separately supersymmetric term* in the Lagrangian density (3.3). The plentiful of these expressions, if not already (3.3) and (3.4), make it clear that 2-dimensional (2,2)-supersymmetric models are far more diverse than $N=1$ supersymmetric models in 4-dimensional spacetime.

Interdependent kinetic terms

In models in higher dimensional spacetimes, the kinetic terms come entirely from the ‘D-terms’ (3.5). In 2-dimensional spacetime, however, we find kinetic terms in all other contributions to the Lagrangian density, L_W in (3.20), L_Σ in (3.21), L_L in (3.23), L_R in (3.26) and L_Q in (3.16). All of these then modify the ‘standard’ (D-term) kinetic terms.

In particular and as discussed already in § 3.4, the λ -terms (3.4f, g), and in fact *all* the nine terms in the Lagrangian density (3.4), contain terms that directly modify the kinetic terms for the components of the various superfields. This type of mechanism was used in Ref. [16] to prove that kinetic terms in fact can be invariant under renormalization flow; indeed, the kinetic terms emerging from all except L_D in (3.4h) are protected by the usual nonrenormalization theorems (see, *e.g.*, p.358 of Ref. [7] for an important caveat).

A minor comment may not be out of place here. Educated on simple models with ‘flat’ kinetic terms such as $\int d^4\zeta \Phi \bar{\Phi} \ni (\partial\phi) \cdot (\partial\bar{\phi})$, one is easy to skim over terms like $(\partial_\pm \ell^{\hat{a}})(\partial_\pm r^{\hat{i}})$ as being trivial. Indeed, the simple D-term $\int d^4\zeta \Lambda \Upsilon$

$$\begin{aligned} & [(\partial_\pm \ell) - 2iL_\pm][(\partial_\pm r) - 2iR_\pm] \\ &= (\partial_\pm \ell)(\partial_\pm r) - 4L_\pm R_\pm - 2i(\partial_\pm \ell)R_\pm - 2iL_\pm(\partial_\pm r) , \\ &= \partial_\pm(\ell \partial_\pm r) - 4L_\pm R_\pm - 2i\partial_\pm(\ell R_\pm) - 2i\partial_\pm(L_\pm r) , \end{aligned} \tag{3.31}$$

turns into a total derivative, except for the non-derivative $L_\pm R_\pm$ term. This clearly carries no dynamics for the spin-0 fields ℓ and r . However, consider $\int d^2\zeta (\bar{D}_+ \Lambda^{\hat{a}})(\bar{D}_- \Upsilon^{\hat{i}}) M_{\hat{a}\hat{i}}(\Phi)$, which is a special case of the third term in (3.9b). Upon fermionic integration, we obtain the ‘kinetic’ terms:

$$-\frac{1}{4}[(\partial_\pm \ell^{\hat{a}}) - 2iL_\pm^{\hat{a}}][(\partial_\pm r^{\hat{i}}) - 2iR_\pm^{\hat{i}}] M_{\hat{a}\hat{i}}(\phi) . \tag{3.32}$$

These definitely do not turn into total derivatives, as long as the ‘metric’ $M_{\hat{a}\hat{i}}(\phi)$ is not constant; in fact, these provide non-trivial ϕ -dependent kinetic (mixing) terms for the oppositely handed bosons $\ell^{\hat{a}}$ and $r^{\hat{i}}$.

Auxiliary fields and (non)linearity

After having expanded the Lagrangian density into component fields by means of fermionic integration, we notice that a number of fields have algebraic equations of motion. By

contrast, fields with differential equations of motion are called physical. For example, consider using the $(\bar{D}_- \Phi^{\bar{\nu}}) M_{\bar{\nu}}$ by itself for a Lagrangian:

$$L_1 = \frac{1}{8} [D_+, \bar{D}_+] D_- (\bar{D}_- \phi^{\bar{\nu}}) M_{\bar{\nu}} + h.c. \quad (3.33)$$

The full component expansion is given in Appendix B.2. From there, we find

$$\begin{aligned} 0 = \frac{\delta L_1}{\delta F^{\bar{\nu}}} &= F^{\mu} (M_{\bar{\nu}, \mu} + \bar{M}_{\mu, \bar{\nu}}) + \psi_-^{\rho} (\psi_+^{\mu} M_{\bar{\nu}, \mu \rho} + \lambda_+^{\hat{a}} M_{\bar{\nu}, \hat{a} \rho}) \\ &\quad - \psi_-^{\rho} (\psi_{L+}^{\tilde{\mu}} \bar{M}_{\rho, \tilde{\mu} \bar{\nu}} + \chi_+^{\tilde{\beta}} \bar{M}_{\rho, \tilde{\beta} \bar{\nu}} + \bar{\lambda}_+^{\hat{a}} \bar{M}_{\rho, \hat{a} \bar{\nu}}) , \end{aligned} \quad (3.34)$$

and the conjugate relation by varying with respect to F^{μ} . These are the equations of motion for F^{μ} and $F^{\bar{\nu}}$. Wherever the hermitian matrix $(M_{\bar{\nu}, \mu} + \bar{M}_{\mu, \bar{\nu}})$ is invertible, the equations of motion can be solved for all the F 's, and these solutions may be substituted back into the Lagrangian density. Writing $M^{\mu \bar{\nu}}$ for the matrix inverse of $(M_{\bar{\nu}, \mu} + \bar{M}_{\mu, \bar{\nu}})$, we'd have

$$F^{\mu} = M^{\mu \bar{\nu}} [\psi_-^{\rho} (\psi_{L+}^{\tilde{\mu}} \bar{M}_{\rho, \tilde{\mu} \bar{\nu}} + \chi_+^{\tilde{\beta}} \bar{M}_{\rho, \tilde{\beta} \bar{\nu}} + \bar{\lambda}_+^{\hat{a}} \bar{M}_{\rho, \hat{a} \bar{\nu}}) - \psi_-^{\rho} (\psi_+^{\sigma} M_{\bar{\nu}, \sigma \rho} + \lambda_+^{\hat{a}} M_{\bar{\nu}, \hat{a} \rho})] , \quad (3.35)$$

and a similar expression for $\Phi^{\bar{\nu}}$. Clearly, this is nonlinear in the fields: roughly, F is quadratic in the fermions, but the details depend on the choice of the Lagrangian density. Its substitution back in the Lagrangian density produces additional cubic and quartic terms, and so affects the interactions between the various fields.

The relation (3.35) also enters in another important place. Prior to elimination of the F 's, the supersymmetry transformation of ψ_- is

$$\delta_{\epsilon} \psi_- = \frac{1}{\sqrt{2}} D_- (\epsilon \cdot Q + \bar{\epsilon} \cdot \bar{Q}) \Phi = -\sqrt{2} \epsilon^+ F + \dots \quad (3.36)$$

which is linear in the fields (including the omitted terms). After substituting (3.35), this transformation rule becomes nonlinear in fields, the details of this nonlinearity depending on the choice of the Lagrangian density. We will refer to these two incarnations of the supersymmetry transformation rules as ‘linear’ and ‘nonlinear’.

It is therefore logically backward to postulate a (nonlinear) set of supersymmetry transformations involving only physical fields, and then seek an invariant Lagrangian density. Although the *linear* supersymmetry transformation rules for a superfield only depend on its definition, *i.e.*, which superconstraints it might satisfy, the *nonlinear* transformation rules depend on the choice of the Lagrangian density through the equations of motion for the auxiliary fields. In particular, modifications of the Lagrangian density (such as through varying moduli) also modify the non-linear supersymmetry transformation rules. In contradistinction, the linear transformation rules, with the auxiliary fields uneliminated, remain unchanged¹⁰⁾, determined entirely by the nature of the superfield itself.

¹⁰⁾ May the erudite Reader forgive this decidedly un-English but accurate agglutination.

4. Explicit Constraining

The Lagrangian density (3.3) has been obtained as a sum of Berezin integrals over suitable subsets of the fermionic coordinates. The superderivative superconstraints (2.1) and (2.3) (and easily also (2.19), should one desire) were *implicitly* and *automatically* used in the process.

Of course, the Reader may instead wish to include such (super)constraints explicitly. To that end, and as disclaimed above, here we examine some simple Lagrangians which enable an explicit inclusion of (super)constraints, and so a more standard treatment of the unidexterous superfields. The present treatment, while inspired by that of Ref. [9], is hopefully a self-sufficient introduction to the topic.

4.1. Chiral bosons

One standard Lagrangian density for a ‘chiral boson’ is [8]:

$$L_{CB} = (\partial_{\mp}\phi)(\partial_{\pm}\phi) + Z_{\mp}^{\pm}(\partial_{\pm}\phi)^2 . \quad (4.1)$$

The inclusion of the constraint through a linear term, $Z^{\pm}(\partial_{\pm}\phi)$ would have left the dynamical degrees of freedom unhalved. Although $\phi(\sigma^{\pm})$ is gauged away, besides $\phi(\sigma^{\pm})$ we also end up with $\tilde{Z}^{\pm}(\sigma^{\pm})$, where $\tilde{Z}^{\pm} \stackrel{\text{def}}{=} (Z^{\pm} + \partial_{\mp}\phi)$, at least classically [17,18]. Note that upon imposing the constraint, $\partial_{\pm}\phi = 0$, the physical field $\phi(\sigma^{\pm})$ and the Lagrange multiplier, $Z_{\mp}^{\pm}$, drop out of the Lagrangian density (4.1), indicating that this Lagrangian density is invariant under a gauge symmetry. A straightforward calculation verifies that (4.1) transforms into a total derivative under the Siegel symmetry transformation:

$$\begin{aligned} \delta_S \phi &= \kappa^{\pm}(\partial_{\pm}\phi) , \\ \delta_S Z_{\mp}^{\pm} &= -(\partial_{\mp}\kappa^{\pm}) - (\partial_{\pm}\kappa^{\pm})Z_{\mp}^{\pm} + \kappa^{\pm}(\partial_{\pm}Z_{\mp}^{\pm}) . \end{aligned} \quad (4.2)$$

The Lagrangian density (4.1) is also invariant with respect to a global inhomogeneous (axial) symmetry:

$$\delta_{\alpha}\phi = \alpha , \quad \delta_{\alpha}Z_{\mp}^{\pm} = 0 , \quad \alpha = \text{const} , \quad (4.3)$$

which allows the interpretation of ϕ as a ‘self-dual gauge scalar’. (Self-duality here means that $\partial_{\pm}\phi = (\sigma_3^{ij} - \epsilon^{ij})\partial_j\phi = 0$, where $i, j = 0, 1$ and $\epsilon^{01} = 1$.)

The following two quite obvious observations will guide the construction of the (2,2)-supersymmetric version:

1. the second term enforces the constraint(s) via the Lagrange multiplier;
2. the remaining part of the Lagrangian (here, the first term) pertains to the unconstrained field.

These two features will then be required of the (2,2)-supersymmetric version.

4.2. Chiral lepton

Consider a simple model, with one chiral superfield, Φ , governed by the following simple Lagrangian density:

$$L_{CL} = \frac{1}{4} \int d^4\zeta \left[\Phi \bar{\Phi} + \frac{1}{2} \mathbf{Z}^= |D_- \Phi|^2 \right]. \quad (4.4)$$

Variation with respect to the Lagrange multiplier (general) superfield, $\mathbf{Z}^=$, then produces the quadratic superconstraint, $|D_- \Phi|^2 = 0$, which restricts the chiral superfield into a chiral lepton superfield (2.3a). This one time we did include a term which ultimately leads to higher derivatives in the Lagrangian density. Defining the components of $\mathbf{Z}^=$ as:

$$Z_{\bar{D}}^- = -\frac{1}{8} (D_- \bar{D}_- \bar{D}_+ D_+ + D_+ \bar{D}_+ \bar{D}_- D_-) \mathbf{Z}^= |, \quad Z^= = \mathbf{Z}^= |, \quad (4.5a)$$

$$Z_{\lambda+}^- = -\frac{1}{4\sqrt{2}} [D_+, \bar{D}_+] \bar{D}_- \mathbf{Z}^= |, \quad Z_{\lambda+}^- = -\frac{1}{4\sqrt{2}} [D_+, \bar{D}_+] D_- \mathbf{Z}^= |, \quad (4.5b)$$

$$Z_{\lambda-}^- = -\frac{1}{4\sqrt{2}} [D_-, \bar{D}_-] \bar{D}_+ \mathbf{Z}^= |, \quad Z_{\lambda-}^- = -\frac{1}{4\sqrt{2}} [D_-, \bar{D}_-] D_+ \mathbf{Z}^= |, \quad (4.5c)$$

$$Z_{\bar{F}}^- = -\frac{1}{2} \bar{D}_+ \bar{D}_- \mathbf{Z}^= |, \quad Z_{\bar{F}}^- = -\frac{1}{2} D_- D_+ \mathbf{Z}^= |, \quad (4.5d)$$

$$Z_{\mp}^- = -\frac{1}{4} [D_+, \bar{D}_+] \mathbf{Z}^= |, \quad Z_{\mp}^- = -\frac{1}{4} [D_-, \bar{D}_-] \mathbf{Z}^= |, \quad (4.5e)$$

$$Z_{\bar{X}}^- = -\frac{1}{2} D_+ \bar{D}_- \mathbf{Z}^= |, \quad Z_{\bar{X}}^- = -\frac{1}{2} D_- \bar{D}_+ \mathbf{Z}^= |, \quad (4.5f)$$

$$Z_{\psi\alpha}^- = -\frac{1}{\sqrt{2}} D_\alpha \mathbf{Z}^= |, \quad Z_{\psi\alpha}^- = -\frac{1}{\sqrt{2}} \bar{D}_\alpha \mathbf{Z}^= |, \quad (4.5g)$$

where $\alpha = -, +$. Note that all sixteen component fields of $\mathbf{Z}^=$ act as Lagrange multipliers.

The simple Lagrangian density (4.4) contains, among many, the following terms:

$$L_{CL} = \frac{1}{2} [(\partial_=\phi)(\partial_+\bar{\phi}) + (\partial_+\phi)(\partial_=\bar{\phi})] - F\bar{F} + \frac{i}{2} [(\psi_- \overleftrightarrow{\partial}_+ \bar{\psi}_-) + (\psi_+ \overleftrightarrow{\partial}_- \bar{\psi}_+)] \quad (4.6a)$$

$$+ Z_{\bar{D}}^- [\psi_- \bar{\psi}_-] \quad (4.6b)$$

$$+ Z_{\mp}^- [i(\partial_=\psi_-) \bar{\psi}_- - (\partial_=\phi)(\partial_=\bar{\phi})] - Z_{\mp}^- [i(\partial_=\psi_-) \bar{\psi}_- - F\bar{F}] \quad (4.6c)$$

$$+ \frac{1}{2} Z^= [(\partial_+\partial_=\psi_-) \bar{\psi}_- + \psi_- (\partial_+\partial_=\bar{\psi}_-) - 2(\partial\psi_+)(\partial\bar{\psi}_+) \\ + i(\partial_+\partial_=\phi)(\partial_=\bar{\phi}) - i(\partial_=\phi)(\partial_+\partial_=\bar{\phi}) + i(\partial_=\bar{F})\bar{F} - iF(\partial_=\bar{F})] \quad (4.6d)$$

+ ...

where $(A \overleftrightarrow{\partial} B) \stackrel{\text{def}}{=} [A(\partial B) - (\partial A)B]$.

Variation with respect to $Z_{\bar{D}}^-$ imposes $|\psi_-|^2 = 0$ (so that $\psi_- = 0 = \bar{\psi}_-$), whereupon the subsequent constraints simplify. Variation with respect to $Z_{\mp}^-$ and $Z_{\mp}^-$ then imposes the constraints $|\partial_=\phi|^2 = 0$ and $|F|^2 = 0$ (so that $\partial_=\phi = 0 = \partial_=\bar{\phi}$ and $F = 0 = \bar{F}$), respectively. This further simplifies the last shown constraint, so that variation with respect to $Z^=$ imposes the constraint $|\partial_=\psi_+|^2 = 0$ (so that $\partial_=\psi_+ = 0$). All the sixteen component fields of $\mathbf{Z}^=$ drop out of the Lagrangian upon enforcing these constraints, indicating the presence of a generalization of the Siegel symmetry (4.2). The net effect of the superconstraint term

in (4.4) then is to impose the constraints (and their conjugates):

$$\psi_- = 0, \quad F = 0; \quad (4.7a)$$

$$\partial_=\phi = 0, \quad \partial_=\psi_+ = 0; \quad (4.7b)$$

The first two of these, Eqs. (4.7a), halve the number of component fields in the chiral superfield (2.1a) to those of the chiral lepton (2.3a); the second two, Eqs. (4.7b), ensure that the remaining component fields are left-movers. This completes the (initial, primary) constraining of the chiral superfield to a lepton chiral; the other quartoid fields (2.3) can be treated in the same fashion, *mutatis mutandis*. Furthermore, a similar treatment of the unidexterous haploid superfields (2.1e, f) seems equally straightforward, albeit rather more beset with technical detail.

The choice (4.4) is not necessarily the most convenient. The simple alternative

$$L_{CL} = \frac{1}{4} \int d^4\zeta \Phi \bar{\Phi} + \frac{1}{4} \int d\zeta^+ \mathbf{Z}_{\mp}^- |D_- \Phi|^2, \quad (4.8)$$

produces a subtly different result. Defining a partial list of component fields of $\mathbf{Z}_{\mp}^-$ as:

$$\mathbf{Z}_{\mp}^- = \mathbf{Z}_{\mp}^-|, \quad Z^- = \frac{1}{4} [D_-, \bar{D}_-] \mathbf{Z}_{\mp}^-|, \quad (4.9a, b)$$

$$\zeta_+^- = \frac{1}{\sqrt{2}} D_- \mathbf{Z}_{\mp}^-|, \quad \zeta_+^- = -\frac{1}{\sqrt{2}} \bar{D}_- \mathbf{Z}_{\mp}^-|, \quad (4.9c, d)$$

the constraint term becomes

$$\begin{aligned} & \frac{1}{8} [D_-, \bar{D}_-] \mathbf{Z}_{\mp}^-| D_- \Phi|^2| \\ &= Z^- |\psi_-|^2 - Z_{\mp}^- [| \partial_=\phi|^2 + \frac{i}{2} (\psi_- \overleftrightarrow{\partial}_= \bar{\psi}_-)] + i \zeta_+^- (\partial_=\phi) \bar{\psi}_- + i \zeta_+^- \psi_- (\partial_=\bar{\phi}). \end{aligned} \quad (4.10)$$

Variation with respect to Z^- annihilates the fermions, $\psi_-, \bar{\psi}_-$. Thereupon, variation with respect to $\mathbf{Z}_{\mp}^-$ sets $\partial_=\phi = 0 = \partial_=\bar{\phi}$. The auxiliary component field F and the fermions $\psi_+, \bar{\psi}_+$ appear only in the expansion of the first ('kinetic') term. It is then their equation of motion that sets $F = 0 = \bar{F}$ and $\partial_=\psi_+ = 0 = \partial_=\bar{\psi}_+$. Thus, we are again left with the component fields of a (quartoid) chiral lepton superfield, and the sixteen components fields of $\mathbf{Z}_{\mp}^-$ which disappear from the action upon enforcing the constraints.

So, whereas the superconstraint term in (4.4) by itself reduces the chiral superfield to a chiral lepton and annihilates the 'kinetic' term in the process, the superconstraint term in (4.8) does only 'half' of that, the rest of the constraining is effected by the equations of motion, derived from the 'kinetic' term in (4.8). Having demonstrated these two possibilities, we leave the application- and taste-dependent choice between these (and possibly other) alternatives to the Reader.

4.3. The general model

The skeptical Reader might inquire as to the relevancy of the above simple model (4.4) to the vastly more complicated Lagrangian density (3.3), which are quite freely interspersed with the sundry unidexterous haploid and quartoid superfields. To that end, note the following simple facts about this Lagrangian density:

1. Terms that would involve any of the constraints (4.7) are *implicitly* projected out of the Lagrangian density (3.3), during the fermionic integration—by its very definition.
2. The Lagrangian density (3.3) does not include any of the Lagrange multiplier superfields enforcing the superconstraints (2.1) and (2.3), because of item 1.
3. The Siegel transformations (4.2) turn unidexterous field(s) into a multiple of the unidexterity superconstraint. That is, the (unidexterity) superconstraints are the generators of the Siegel symmetry.

The Lagrangian density (3.3), being *implicitly* restricted to the constrained (super)field space, is then trivially an invariant of the Siegel symmetry. So, at least at the classical level, there should be no obstacle to using the unidexterous superfields in all the ways they appear herein.

One could instead relax the unidexterity superconstraints (2.1e, f) on the Λ 's and Υ 's, and impose these superconstraints explicitly through adding suitable Lagrangian multiplier terms to (a suitable subset of) the Lagrangian density (3.3). Finding thereupon a suitable generalization of the Siegel symmetry for this modified Lagrangian density, should guarantee a well-defined interactive quantum theory. This, or any other of the many approaches to constrained quantization, is however beyond our present scope. The Reader weary of the unidexterous quantization issues may of course safely use the Lagrangian density (3.3) with the unidexterous superfields carefully extracted.

On the other hand, the axial symmetry analogue of (4.3) for each of the unidexterous superfields is explicitly broken by each term where an inhomogeneously transforming component field of a unidexterous superfield appears without a spacetime derivative acting on it. Many, however, do not break the axial symmetries: consider the term $\int d^2\varsigma (\bar{D}_+ \Lambda^{\hat{a}})(\bar{D}_- \Upsilon^{\hat{i}}) M_{\hat{a}\hat{i}}(\Phi)$, obtained as a special case from (3.9b), which upon fermionic integration yields

$$\begin{aligned}
& -\frac{1}{4}[(\partial_{\neq} \ell^{\hat{a}}) - 2iL_{\neq}^{\hat{a}}][(\partial_{=} r^{\hat{i}}) - 2iR_{=}^{\hat{i}}] M_{\hat{a}\hat{i}}(\phi) + \psi_{-}^{\mu} \psi_{+}^{\nu} \lambda_{+}^{\hat{a}} \varrho_{-}^{\hat{i}} M_{\hat{a}\hat{i},\mu\nu}(\phi) \\
& + \{ F^{\mu} \lambda_{+}^{\hat{a}} \varrho_{-}^{\hat{i}} + \frac{i}{2} \psi_{-}^{\mu} [(\partial_{\neq} \ell^{\hat{a}}) - 2iL_{\neq}^{\hat{a}}] \varrho_{-}^{\hat{i}} + \frac{i}{2} \psi_{-}^{\mu} \lambda_{+}^{\hat{a}} [(\partial_{=} r^{\hat{i}}) - 2iR_{=}^{\hat{i}}] \} M_{\hat{a}\hat{i},\mu}(\phi) .
\end{aligned} \tag{4.11}$$

This involves $\ell^{\hat{a}}$ and $r^{\hat{i}}$, the lowest component fields of the lefton and righton superfields, only through their spacetime derivatives. Consequently, it is invariant under the two inhomogeneous ‘axial’ transformations: $\delta_{\alpha} \ell^{\hat{a}} = \alpha^{\hat{a}}$ (where $\partial_{\neq} \alpha^{\hat{a}} = 0$), and $\delta_{\alpha} r^{\hat{i}} = \alpha^{\hat{i}}$ (where $\partial_{=} \alpha^{\hat{i}} = 0$), with respect to which all other fields are held inert.

The inquisitive Reader is invited to sift through the multitude of individually supersymmetric parts of the Lagrangian density (3.3) and identify all the terms with such property, should this be a requirement of a desired application.

Finally, the constraining Lagrangian density (4.4) could serve as a prototype of an *explicit* constraining of the various constrained superfields [8]. For example, given an unconstrained complex (2,2)-superfield, Ω , and two unconstrained Lagrange multiplier superfields, $\mathbf{Z}^{-}$ and $\mathbf{Z}^{+}$, the Lagrangian term

$$\frac{1}{8} \int d^4\varsigma \left[\mathbf{Z}^{-} (\bar{D}_{-} \Omega) (D_{-} \bar{\Omega}) + \mathbf{Z}^{+} (\bar{D}_{+} \Omega) (D_{+} \bar{\Omega}) \right] , \tag{4.12}$$

will constrain Ω into a chiral superfield. Again, the maintenance of the generalization of the Siegel symmetry off the constrained superfield subspace and quantum issues are beyond our present scope.

5. Wave-Function(al)s and Geometry

The Hilbert space of supersymmetric field theories is in many regards much simpler than the one in models without supersymmetry [19,20,21,22,23,24]. Although many of the known results have been derived for 2-dimensional supersymmetric field theories, the lack of present generality warrants another look at these issues.

5.1. Vacua and Wave-Function(al)s

The wave-function(al)s in supersymmetric theories are functions of the superfields and so admit a formal expansion in the fermionic fields. Local such wave function(al)s (in the 2-dimensional spacetime sense) will then have a terminating Taylor series owing to the finite number of fermionic fields and their anticommutivity, and so must be multinomials in fermionic fields.

A hallmark of fermionic fields is the fact that their kinetic terms are first order in time derivatives, so half the fermionic fields turn out to be conjugate momenta to the other half. The exact pairing of course depends on the exact choice of the Lagrangian density (3.3). On the other hand, the identification of the ‘canonical coordinate’ and the ‘canonical momentum’ within any one canonically conjugate pair of fermion fields is completely arbitrary.

In a second-quantized theory, we define a (Dirac) ‘vacuum’ state, $|0\rangle$, which is annihilated by precisely one half of the fermionic fields. Except, of course, the choice of precisely which half will annihilate $|0\rangle$ remains free. As well known, Eqs. (1.2) imply

$$\{Q_-, \bar{Q}_-\} + \{Q_+, \bar{Q}_+\} = 4H, \quad (5.1)$$

and so

$$\begin{aligned} \langle H \rangle &= \frac{1}{4} \langle \{Q_-, \bar{Q}_-\} \rangle + \frac{1}{4} \langle \{Q_+, \bar{Q}_+\} \rangle \\ &= \frac{1}{4} [|Q_-|^2 + |\bar{Q}_-|^2 + |Q_+|^2 + |\bar{Q}_+|^2] \geq 0, \end{aligned} \quad (5.2)$$

the equality reached precisely for supersymmetric states, *i.e.*, those annihilated by all four supercharges. That is, in supersymmetric models, the Hamiltonian is positive definite, and its global minima (of zero energy, hence *the true vacua*) are the supersymmetric states.

It is then the supersymmetric states, *the true vacua*, that we seek in a supersymmetric model, so as to erect the full Hilbert space by acting on these vacua with creation operators. In turn, the vacua, *i.e.*, supersymmetric states are easily constructed from the ‘vacua’ $|0\rangle$.

Single species models

Even if considering just a single species of fermionic fields, such as $\psi_\pm^\mu, \psi_\pm^{\bar{\nu}}$ which appear in chiral superfields and their conjugates (as used in Wess-Zumino model), respectively, one easily finds six *logically possible* choices of such a $|0\rangle$:

$$(\mathbf{c}, \mathbf{c}) \text{ vacuum : } \quad \psi_+^\mu |0\rangle_{(c,c)} = 0 = \psi_-^\mu |0\rangle_{(c,c)} ; \quad (5.3a)$$

$$(\mathbf{a}, \mathbf{a}) \text{ vacuum : } \quad \psi_+^{\bar{\mu}} |0\rangle_{(a,a)} = 0 = \psi_-^{\bar{\mu}} |0\rangle_{(a,a)} ; \quad (5.3b)$$

$$(\mathbf{a}, \mathbf{c}) \text{ vacuum : } \quad \psi_+^{\bar{\mu}} |0\rangle_{(a,c)} = 0 = \psi_-^\mu |0\rangle_{(a,c)} ; \quad (5.3c)$$

$$(\mathbf{c}, \mathbf{a}) \text{ vacuum : } \quad \psi_+^\mu |0\rangle_{(c,a)} = 0 = \psi_-^{\bar{\mu}} |0\rangle_{(c,a)} ; \quad (5.3d)$$

$$\text{Left vacuum : } \quad \psi_-^\mu |0\rangle_L = 0 = \psi_-^{\bar{\mu}} |0\rangle_L . \quad (5.3e)$$

$$\text{Right vacuum : } \quad \psi_+^\mu |0\rangle_R = 0 = \psi_+^{\bar{\mu}} |0\rangle_R . \quad (5.3f)$$

We have assumed here that a degree of uniformity is to be required of the chiral superfields and their conjugates, so that all (un)conjugate fermions appear in the defining equations (5.3) on the same footing. In particular, the definitions (5.3) are invariant with respect to any invertible linear transformation among the chiral superfields Φ^μ (which commutes with the Lorentz symmetry, and so preserves spin), accompanied by the conjugate transformation among their antichiral conjugates, $\Phi^{\bar{\mu}}$. Clearly, if (global) invariance is to be required only with respect to a subset of these transformations, several additional choices of $|0\rangle$ will be possible.

For any one choice of $|0\rangle$, the half of the fermionic fields which do not annihilate $|0\rangle$ then act as creation operators, and states are constructed as multinomials in these ‘creation’ fermions, acting on $|0\rangle$. Naturally, these wave function(al)s (states) built upon these ‘vacua’ will be called by the same names: that is, a (c, c) state will be built from a (c, c) vacuum, $|0\rangle_{(c,c)}$, an (a, c) state from $|0\rangle_{(a,c)}$, and so on.

Now, the kinetic term for the fermions is typically something like

$$\begin{aligned} L_{\text{kin}}^{(F)} &= iG_{\mu\bar{\nu}} [\psi_+^\mu (\partial_- \psi_+^{\bar{\nu}}) + \psi_-^\mu (\partial_+ \psi_-^{\bar{\nu}})] , \\ &= iG_{\mu\bar{\nu}} (\psi_+^\mu \dot{\psi}_+^{\bar{\nu}} + \psi_-^\mu \dot{\psi}_-^{\bar{\nu}} - \psi_+^\mu \psi_+^{\bar{\nu}} + \psi_-^\mu \psi_-^{\bar{\nu}}) , \end{aligned} \quad (5.4)$$

where $\dot{\psi}_\pm \stackrel{\text{def}}{=} \partial_0 \psi_\pm$ is the time-derivative of $\psi_\pm$, $\dot{\psi}_\pm \stackrel{\text{def}}{=} \partial_1 \psi_\pm$ its space-derivative, and $G_{\mu\bar{\nu}}$ is the metric determined by the choice of the Lagrangian density (3.3). Therefore, $i\psi_{\pm\mu} \stackrel{\text{def}}{=} iG_{\mu\bar{\nu}} \psi_\pm^{\bar{\nu}}$ is the canonical conjugate of $\psi_\pm^\mu$, so that

$$\{i\psi_{\pm\mu}, \psi_\pm^\nu\} = i\delta_\mu^\nu, \quad i.e. \quad \{\psi_\pm^{\bar{\nu}}, \psi_\pm^\mu\} = G^{\mu\bar{\nu}} . \quad (5.5)$$

Notice now that the fifth and sixth logical possibilities, tentatively called the ‘Left’ and ‘Right’ vacua in fact must vanish, since the two conditions (5.3e) may be combined to yield

$$0 = \{\psi_-^\mu, \psi_-^{\bar{\nu}}\} |0\rangle_L = -iG^{\mu\bar{\nu}} |0\rangle_L , \quad (5.6)$$

whereupon $|0\rangle_L \equiv 0$ is inevitable. So, unlike (2,2)-superfields in 2-dimensional spacetime, there are only four types (two conjugate pairs) of (2,2)-vacua — for a single species of fermions.

In a σ -model, the bosonic coordinates $\phi^\mu, \phi^{\bar{\nu}}$ are identified as target space coordinate-valued world-sheet scalars. That is, they map the world-sheet (*domain*) into X (the *target space*). In a Fourier expansion (locally in X), the 0-wavevector modes may be identified with local coordinates $z, \bar{z}$ in X . Then, half of the fermions $\psi_\pm, \bar{\psi}_\pm$ can be identified with the differentials $dz, d\bar{z}$, and the other half with the partials $\partial, \bar{\partial}$. We adopt:

$$\begin{aligned} G_{\mu\bar{\nu}}\psi_-^\mu &\mapsto \partial_{\bar{\nu}} , & \psi_-^{\bar{\nu}} &\mapsto dz^{\bar{\nu}} , \\ \psi_+^\mu &\mapsto dz^\mu , & G_{\mu\bar{\nu}}\psi_+^{\bar{\nu}} &\mapsto \partial_\mu , \end{aligned} \tag{5.7}$$

That the anticommutivity of ψ_+^μ and $\psi_-^{\bar{\nu}}$ matches the anticommutivity of $dz, d\bar{z}$, standard in differential geometry. However, whereas the ‘momenta’ $G_{\mu\bar{\nu}}\psi_-^\mu$ and $G_{\mu\bar{\nu}}\psi_+^{\bar{\nu}}$ also anticommute, the basis elements of the tangent bundle, $\partial_{\bar{\nu}}$ and ∂_μ *commute*. Nevertheless, the above identification has proven of great use in the analysis of the geometry of supersymmetric field theory models.

With the identifications (5.7), the (c, c) wave function

$$|p, q; h\rangle_{(c, c)} = h_{\bar{\nu}_1 \dots \bar{\nu}_q}^{\mu_1 \dots \mu_p}(\phi, \bar{\phi}) \psi_{+\mu_1} \dots \psi_{+\mu_p} \psi_{-}^{\bar{\nu}_1} \dots \psi_{-}^{\bar{\nu}_q} |0\rangle_{(c, c)} \tag{5.8}$$

corresponds to the $\wedge^p T_X$ -valued q -form

$$h_{\bar{\nu}_1 \dots \bar{\nu}_q}^{\mu_1 \dots \mu_p}(\phi, \bar{\phi}) \partial_{\mu_1} \wedge \dots \wedge \partial_{\mu_p} \wedge dz^{\bar{\nu}_1} \wedge \dots \wedge dz^{\bar{\nu}_q} .$$

Similarly, the (a, c) wave function

$$|p, q; \omega\rangle_{(a, c)} = \omega_{\mu_1 \dots \mu_p \bar{\nu}_1 \dots \bar{\nu}_q}(\phi, \bar{\phi}) \psi_+^{\mu_1} \dots \psi_+^{\mu_p} \psi_{-}^{\bar{\nu}_1} \dots \psi_{-}^{\bar{\nu}_q} |0\rangle_{(a, c)} \tag{5.9}$$

corresponds to the $\wedge^p T_X^*$ -valued q -form

$$\omega_{\mu_1 \dots \mu_p \bar{\nu}_1 \dots \bar{\nu}_q}(\phi, \bar{\phi}) dz^{\mu_1} \wedge \dots \wedge dz^{\mu_p} \wedge dz^{\bar{\nu}_1} \wedge \dots \wedge dz^{\bar{\nu}_q} .$$

Supersymmetric states

States which are invariant with respect to supersymmetry transformations are of very special interest [19]; we recall a few important facts here. Representing a supersymmetric state, let $|*\rangle$ stand for a wave-function annihilated by all supercharges. Then

$$\{ Q_\pm, \bar{Q}_\pm \} |*\rangle = 0 \quad \Leftrightarrow \quad (H \pm p) |*\rangle = 0 , \tag{5.10}$$

which is equivalent to

$$p |*\rangle = 0 = H |*\rangle . \tag{5.11}$$

That is, translationally invariant states of zero energy are supersymmetric, and *vice versa*.

Now, the precise form of $H, p, Q_\pm$ and $\bar{Q}_\pm$ (as operators on the field space) depends on the precise choice of the Lagrangian density (3.3). Restricting to translationally invariant (world-sheet space-independent) states, is in many ways equivalent to completely disregarding the spatial extension of the world-sheet, and reducing from the 2-dimensional

(supersymmetric) field theory to (supersymmetric) quantum mechanics. In practice, this allows us to *ignore* the momentum density p and all the terms which involve world-sheet spatial derivatives; H correspondingly simplifies to the Fokker-Planck type Hamiltonian. For the case of Wess-Zumino models (those containing only n chiral superfields and their conjugates), these operators have been identified many years ago [21,22].

For the more general case involving more than just one species of superfields (2.1), (2.3), (2.19), *etc.*, this remains unexplored, albeit straightforward.

Three remarks

Before we turn to a cursory look at multiple-species models, three general remarks are in order. These are based on the fact that the adjoint pair of (Dolbeault) differential operators $\bar{\partial} \stackrel{\text{def}}{=} dz^\mu \partial_\mu \wedge$ and $\bar{\partial}^*$, with the only non-zero anticommutation relation $\{\bar{\partial}, \bar{\partial}^*\} = \Delta_{\bar{\partial}}$, where $\Delta_{\bar{\partial}}$ is the $\bar{\partial}$ -Laplacian. Indeed, the correspondence between supersymmetric and cohomology theories, first noted in Ref. [19], stems from the (formal) isomorphism between these and the pair of anticommuting differential operators $D_\pm, \bar{D}_\pm$ with the only non-zero anticommutation relation (1.6).

1. Corresponding to the two pairs ∂, ∂^* , and $\bar{\partial}, \bar{\partial}^*$ in differential geometry, we have $Q_-, \bar{Q}_-$ and $Q_+, \bar{Q}_+$. Then, the supersymmetric analogue of complex conjugation in differential geometry is not hermitian conjugation but *parity*, $\mathbf{P}$. Of course, *upon Euclideanization*, parity does act as complex conjugation on the (bosonic ‘body’ of the) world-sheet Riemann surface. On the other hand, the differential-geometric analogue of hermitian conjugation in supersymmetric models is the formal exchange $\bar{\partial} \leftrightarrow \bar{\partial}^*$, not complex conjugation.
2. The analogue of $\Delta_{\bar{\partial}}$ is $2(H + p)$, *i.e.*, $\partial_\mp$; the analogue of Δ_{∂} is $2(H - p)$, *i.e.*, ∂_+ . When complex conjugation is a symmetry of the target space (as it happens, but not exclusively, on Kähler manifolds), $\Delta_{\bar{\partial}} = \Delta_{\partial}$. The analogous situation with the supersymmetry algebra and its representations happens when $2(H + p) = 2(H - p)$: in the sector of zero-momentum states, where $p|*\rangle = 0$. Equivalently, this implies $\partial_\mp = \partial_+$, which holds on σ^1 -independent functions, *i.e.*, upon dimensional reduction to (supersymmetric) quantum mechanics.
3. Being annihilated by both D_+ and $\bar{D}_+$, the righton superfields, Υ (2.1f), are isomorphic to $\bar{\partial}$ -closed and co-closed forms, that is, to $\bar{\partial}$ -harmonic forms. Similarly, the lefton superfields, Λ (2.1f*e*), are isomorphic to ∂ -closed and co-closed forms, that is, to ∂ -harmonic forms. This correspondence has, to the best of my knowledge, not been observed before and its utility remains unexplored.

Several species models

With more types of superfields, and so more species of fermions, the number of different types (sectors) of ‘vacua’ and wave-function(al)s increases very quickly.

Even if restricting to just the ‘minimal’ haploid superfields, there are many more species of fermions: $\psi_\pm^\mu, \bar{\psi}_\pm^\mu$ from (anti)chiral superfields, $\xi_-^\alpha, \chi_+^\alpha$ and $\xi_-^{\bar{\beta}}, \chi_+^{\bar{\beta}}$ from twisted-(anti)chiral superfields, $\lambda_+^a, \bar{\lambda}_+^a$ (and, if complex, their conjugates) from lefton superfields

and ρ_-^i, ϱ_-^i (and, if complex, their conjugates) from righton superfields. Half of these are canonically conjugate variables to the other half (the precise pairing depending on the precise choice of the Lagrangian density), so we in general expect six types of fermionic ‘creation’ operators, not just two as above. Clearly, there will also be many more choices of $|0\rangle$.

Consider a simple toy example with a single ‘minimal’ haploid superfield of each kind (2.1) and the Lagrangian density

$$L_{\text{toy}} = \frac{1}{4} \int d^4\zeta (\Phi\bar{\Phi} - \Xi\bar{\Xi}) \quad (5.12a)$$

$$+ \frac{1}{4} \left[\int d\zeta \Lambda^\pm (\bar{D}_+ D_+ \Lambda^\pm) + h.c. \right], \quad (5.12b)$$

$$+ \frac{1}{4} \left[\int d\zeta \Upsilon^\pm (\bar{D}_- D_- \Upsilon^\pm) + h.c. \right], \quad (5.12c)$$

where, for simplicity, we require $\Lambda^\pm, \Upsilon^\pm$ to be real. Then, the first two terms produce the standard kinetic terms for the scalar fields ϕ and x , their fermionic superpartners, $\psi_\pm$ and ξ_-, χ_+ , respectively, the auxiliary fields F, X , and their hermitian conjugates. The scalars have their usual canonically conjugate momenta, and ψ_-, ψ_+ and ξ_-, χ_+ are canonically conjugate pairs.

We then focus on the second part (5.12b), which produces (up to total derivatives):

$$L_{\text{toy}}^{(\Lambda)} = \frac{1}{2} [(\partial_\mp \ell^\pm) + 2iL_\mp^\pm]^2 - i(\lambda_+^\pm \overleftrightarrow{\partial}_\mp \lambda_+^\pm). \quad (5.13)$$

The equations of motion for $L_\mp^\pm, \lambda_+^\pm$ and $\bar{\lambda}_+^\pm$, respectively, set:

$$L_\mp^\pm = \frac{i}{2}(\partial_\mp \ell^\pm), \quad \partial_\mp \lambda_+^\pm = 0, \quad \partial_\mp \bar{\lambda}_+^\pm = 0. \quad (5.14)$$

Upon substituting the first of these back into the Lagrangian density, both $L_\mp^\pm$ and $\ell^\pm$ disappear from it, indicating a gauge symmetry. As for the fermions, *on shell*, $\lambda_+^\pm, \bar{\lambda}_+^\pm = \text{const.}$ (Recall that $\partial_- \Lambda^\pm = 0$.)

More importantly, however, the boson $\ell^\pm$ has a well-defined canonically conjugate ($L_\mp^\pm$ -dependent) momentum, and $\lambda_+^\pm$ and $\bar{\lambda}_+^\pm$ are canonically conjugate to each other. Similarly, the last part (5.12c) dictates that $\rho_-^\pm$ and $\varrho_-^\pm$ are canonically conjugate to each other.

Thus, we can define *sixty-four* different (Dirac) vacua $|0\rangle$, annihilated by a choice of six fermions:

$$\left\{ \begin{matrix} \psi_- \\ \bar{\psi}_- \end{matrix} \right\}, \left\{ \begin{matrix} \psi_+ \\ \bar{\psi}_+ \end{matrix} \right\}, \left\{ \begin{matrix} \xi_- \\ \bar{\xi}_- \end{matrix} \right\}, \left\{ \begin{matrix} \chi_+ \\ \bar{\chi}_+ \end{matrix} \right\}, \left\{ \begin{matrix} \lambda_+^\pm \\ \bar{\lambda}_+^\pm \end{matrix} \right\}, \left\{ \begin{matrix} \rho_-^\pm \\ \varrho_-^\pm \end{matrix} \right\}, \quad (5.15)$$

where each braced pair indicates a free choice. Building upon these vacua, there are sixty-four (super)sectors of the Hilbert space—vastly generalizing the (c, c) , (a, c) , (c, a) and (a, a) sectors familiar from Landau-Ginzburg orbifolds, but present in any (2,2)-supersymmetric theory.

5.2. Haploid correlation functions

In a model with the Lagrangian density (3.3), it is easy to define correlation functions for the various terms in (3.3). For example, let us denote by $\delta_i W$ the i^{th} term that can be added to the superpotential W in (3.3b). This means that each $\delta_i W$ is a chiral superfield, and so satisfies Eqs. (2.1a). Then, the correlation function

$$\langle 0 | \delta_1 W \cdots \delta_n W | 0 \rangle \quad (5.16)$$

is also called chiral, since the product $\delta_1 W \cdots \delta_n W$ is chiral. Similarly, a twisted-chiral correlation function is the correlation function of a (product of) twisted-chiral object(s), a lepton correlation function of a (product of) lepton object(s), *etc.*

The study of correlation functions simplifies on noting that in theories with unbroken supersymmetry, the vacua must be invariant under supersymmetry, and so annihilated by the supercharges $Q_\pm, \bar{Q}_\pm$. This induces a number of ‘topological’ results [23,24,25], which will herein be referred to as ‘rigidity’, not unrelated to the mathematically standard notion of ‘rigidity’, and to avoid confusion with the mathematically standard notion of ‘topological’.

The ‘rigidity’ results of the work on chiral correlation functions in Wess-Zumino models [26,27,23,24] has a direct analogue for correlation functions of ambixterous haploid (and all quartoid) superfield operators. Given any two chiral superfields, $\Phi(\sigma_1^\pm, \sigma_1^\mp)$ and $\Phi(\sigma_2^\pm, \sigma_2^\mp)$, which depend on two different points on the world-sheet, $\sigma_i^\pm, \sigma_i^\mp$ for $i = 1, 2$, the basic result is as follows:

$$\begin{aligned} \frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Phi_1 \Phi_2 | 0 \rangle &= \frac{1}{2i} \langle 0 | \{ \bar{Q}_-, (Q_- \Phi_1) \} \Phi_2 | 0 \rangle \\ &= \frac{1}{2i} \underbrace{\langle 0 | \bar{Q}_- (Q_- \Phi_1) \Phi_2 | 0 \rangle}_{=0} + \frac{1}{2i} \langle 0 | (Q_- \Phi_1) \Phi_2 \underbrace{\bar{Q}_- | 0 \rangle}_{=0} = 0, \end{aligned} \quad (5.17a)$$

since $[\bar{Q}_\pm, \Phi_i] = 0$, and

$$\frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Phi_1 \Phi_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ \bar{Q}_+, (Q_+ \Phi_1) \} \Phi_2 | 0 \rangle = 0, \quad (5.17b)$$

in a similar fashion. The same, upon $Q_\pm \leftrightarrow \bar{Q}_\pm$, is of course true for $\langle 0 | \bar{\Phi}_1 \bar{\Phi}_2 | 0 \rangle$.

By the same token,

$$\frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Xi_1 \Xi_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ \bar{Q}_-, (Q_- \Xi_1) \} \Xi_2 | 0 \rangle = 0, \quad (5.17c)$$

$$\frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Xi_1 \Xi_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ Q_+, (\bar{Q}_+ \Xi_1) \} \Xi_2 | 0 \rangle = 0, \quad (5.17d)$$

and the same for $\langle 0 | \bar{\Xi}_1 \bar{\Xi}_2 | 0 \rangle$. The same rigidity will hold for the quartoid superfields (2.3), as they all satisfy the superconstraints of some ambidexterous haploid superfield. This can be remembered easily by regarding Φ_L and Φ_R as two ‘halves’ of Φ , and $\bar{\Phi}_L$ and $\bar{\Phi}_R$ as two ‘halves’ of $\bar{\Phi}$.

The situation is quite different for the unidexterous fields:

$$\frac{\partial}{\partial \sigma_1^-} \langle 0 | \Lambda_1 \Lambda_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ \bar{Q}_-, \underbrace{(Q_- \Lambda_1)}_{=0} \} \Lambda_2 | 0 \rangle = 0 , \quad (5.17b)$$

holds again, but now

$$\frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Lambda_1 \Lambda_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ \bar{Q}_+, (Q_+ \Lambda_1) \} \Lambda_2 | 0 \rangle = \frac{1}{2i} \langle 0 | (Q_+ \Lambda_1) (\bar{Q}_- \Lambda_2) | 0 \rangle \neq 0 . \quad (5.17f)$$

Similarly,

$$\frac{\partial}{\partial \sigma_1^-} \langle 0 | \Upsilon_1 \Upsilon_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ \bar{Q}_-, (Q_- \Upsilon_1) \} \Upsilon_2 | 0 \rangle \neq 0 , \quad (5.17h)$$

$$\frac{\partial}{\partial \sigma_1^\mp} \langle 0 | \Upsilon_1 \Upsilon_2 | 0 \rangle = \frac{1}{2i} \langle 0 | \{ Q_+, (\bar{Q}_+ \Upsilon_1) \} \Upsilon_2 | 0 \rangle = 0 . \quad (5.17h)$$

That is:

1. Corellation functions of (products of) any one type of ambidexterous haploid superfields (2.1a–d) (and any quartoid superfields (2.3) that obey the same superconstraint) are independent of the superfields' positions on the world-sheet.
2. Corellation functions of (products of) unidexterous haploid superfields (2.1e, f) are correspondingly unidexterous functions of the superfields' positions on the world-sheet.

It should be clear that only the constrained superfields defined by the simple superconstraints (2.1) and (2.3), which are linear in both superderivatives and superfields, possess this remarkable property of ‘rigidity’.

5.3. Target space geometry

The field (target) space for our model is spanned by the physical component fields in the collection of superfields $\Phi, \bar{\Phi}, \Xi, \bar{\Xi}, \Lambda, \Upsilon$. The entire Lagrangian density (3.3) depends on a grand total of 3 rank-0, 24 rank-1, 32 rank-2, 12 rank-3 and 6 rank-4 (tensor-like) ‘coefficient functions’ (and their complex conjugates) defined in (3.18), (3.22), (3.24), (3.9) and (3.5). Even if the unidexterous superfields, $\Lambda, \Upsilon, \Phi_L, \Phi_R, \bar{\Phi}_L, \bar{\Phi}_R$ and the non-minimal superfields, $\Theta, \bar{\Theta}, \Pi, \bar{\Pi}$, were to be dropped, the Lagrangian density depends on 3 rank-0 and 4 rank-1 coefficient functions, defined in (3.18), (3.22), (3.9) and (3.5):

$$\begin{aligned} & K(\phi, \bar{\phi}, x, \bar{x}) , \quad W(\phi) , \quad \Sigma(x) , \\ & M_{\bar{\nu}}(\phi, x) , \quad M_{\bar{\beta}}(\phi, x) , \quad \widetilde{M}_{\bar{\nu}}(\phi, \bar{x}) , \quad \widetilde{M}_{\alpha}(\phi, \bar{x}) , \end{aligned} \quad (5.18)$$

and the conjugates of all but K , which is real. Each term from Eqs. (3.18), (3.22), (3.9) and (3.5) featuring one of these functions produces, upon the appropriate fermionic integration, a *separately* and *manifestly supersymmetric* term in the Lagrangian (3.3). Therefore, supersymmetry imposes no further condition on the coefficient functions (5.18) and their complement, as defined in (3.18), (3.22), (3.9) and (3.5).

Recall that the last four displayed coefficient functions in (5.18) contribute both to a variation in the D-term (*i.e.*, a change of K) and a ‘topological’ term such as (3.15). Four other terms,

$$\begin{aligned} & \frac{1}{8}[D_+, \bar{D}_+]D_-[(D_- \Phi^\mu)M_\mu^L + (D_- \Xi^\alpha)M_\alpha^L] + h.c. \\ \text{and } & \frac{1}{8}[D_-, \bar{D}_-]D_+[(D_+ \Phi^\mu)\widetilde{M}_\mu^R + (D_+ \Xi^{\bar{\beta}})\widetilde{M}_{\bar{\beta}}^R] + h.c. \end{aligned} \quad (5.19)$$

vanish on retaining only the chiral and twisted-chiral superfields: the coefficient functions M_μ^L, M_μ^R become constants, whereupon the repeated superderivative annihilates the term.

Clearly, these rank-0...4, ‘coefficient functions’ are rank-0...4 tensor-like objects over the field space. However, it must be emphasized that these *need not be global tensors* of such rank! Consider for example the term $(\bar{D}_- \Phi^{\bar{\nu}})M_{\bar{\nu}}(\bar{\Phi}_L, \Phi, \Xi, \Lambda)$ from Eq. (3.9b). The (co)vector $M_{\bar{\nu}}$ need not be a global (co)vector field over the space coordinatized by the (lowest components of) $\Phi^\mu, \Phi^{\bar{\nu}}$. In particular, as used in Eq. (3.11) and below, the (co)vector $M_{\bar{\nu}}$ may well be a contraction

$$\Phi^\mu M_{\mu\bar{\nu}} + \Xi^\alpha M_{\alpha\bar{\nu}} + \Lambda^{\hat{a}} M_{\hat{a}\bar{\nu}} + \dots \quad (5.20)$$

Thus, the fact that in some application, the space coordinatized by the (lowest components of) $\Phi^\mu, \Phi^{\bar{\nu}}$ does not admit a global 1-form $M_{\bar{\nu}} d\phi^{\bar{\nu}}$ does not preclude the inclusion in the Lagrangian density of the first λ -term in Eq. (3.9b).

Furthermore, the coefficient functions in (3.18), (3.22), (3.9) and (3.5) all admit some degree of arbitrariness, generalizing the well-known ‘Kähler symmetry’, $K \simeq K + f(\Phi) + \bar{f}(\bar{\Phi})$, where $f(\Phi)$ is an arbitrary function of the chiral superfields, and $\bar{f}$ its conjugate. In point of fact, the function K in (3.5) is defined only up to the very general reparametrization:

$$K(\Phi, \bar{\Phi}, \Xi, \dots) \simeq K + f_{(-)} + f_{(+)} + \bar{f}_{(-)} + \bar{f}_{(+)} , \quad (5.21)$$

where

$$\bar{D}_- f_{(-)} = 0 = \bar{D}_+ f_{(+)} , \quad D_- \bar{f}_{(-)} = 0 = D_+ \bar{f}_{(+)} . \quad (5.22)$$

That is, $f_{(\pm)}$ and $\bar{f}_{(\pm)}$ are *arbitrary* functions of any of the (superderivative) superfields annihilated by any one of the superderivatives; see Eqs. (2.30). For the other coefficient functions, the reparametrizing functions (such as the f ’s above) must be annihilated by at least one more superderivative than the coefficient function that they reparametrize. For example, the superpotential obeys

$$W \simeq W + g_L + g_R , \quad (5.23)$$

where g_L, g_R are a chiral lefton and a chiral righton function. Clearly, g_L, g_R are nonzero only in models that involve chiral leftons and chiral rightons.

It is then patently clear that the geometry of the field (target) space of general (2,2)-supersymmetric models is considerably more general than just Kähler manifolds with an assortment of (holomorphic) vector bundles (or even sheaves). As it was known since Ref. [1], even a model depending on just the first one of the seven functions (5.18) provides more general geometries, further augmented by the two ‘superpotentials’, $W(\phi)$ and $\Sigma(\phi)$.

The continued extension of the geometrical structure relying on the inclusion of rank-2 coefficient functions $W_{\alpha\bar{\beta}}$ and $\Sigma_{\mu\bar{\nu}}$ was noted in Ref. [4]. These appear through the F-terms

$$\frac{1}{2} \int d^2\zeta (\bar{D}_+\Xi^\alpha)(\bar{D}_-\Xi^{\bar{\beta}})W_{\alpha\bar{\beta}}(\Phi) , \quad \text{and} \quad \frac{1}{2} \int d^2\zeta (D_+\Phi^\mu)(\bar{D}_-\Phi^{\bar{\nu}})\Sigma_{\mu\bar{\nu}}(\Xi) , \quad (5.24)$$

which are in fact special cases of the first two λ -terms (3.9b). To the best of my knowledge, this is the first time the four 1-forms in (5.18) have made their public appearance. Of course, if one insists on restricting to only chiral, twisted-chiral superfields and their conjugates, the seven objects (5.18) admittedly exhausts the list (subject to our general restrictions), with the latter four ‘merely’ modifying the D-terms and contributing ‘topological’ terms such as (3.15).

Note that, except for K which is real, all the coefficient functions are analytic of their arguments, and so holomorphic functions of their complex arguments. The Lagrangian terms in which they appear are not $\int d^4\zeta$ -integrated terms, whence they fall under the scope of the non-renormalization theorems [6,7]. More precisely, their contribution to the D-terms is subject to renormalization, but the complement of these contributions is not. That is, the Lagrangian terms involving all but the first of the functions (5.18) (and all those omitted in (5.18)!) are expected to provide marginal operators (and so deformations) in a quantum theory.

6. Summary, Outlook and Conclusions

In an attempt to provide a comprehensive but comprehensible intrinsic study of (2,2)-supersymmetric models in 2-dimensional spacetime, we purposefully avoid constructions based on dimensional reduction from 4-dimensions.

A self-contained discussion of types of constrained superfields was given in § 2. These include *haploid* (*quartoid*) superfields, which depend on a half (quarter) of the fermionic coordinates. The ‘minimal’ haploid superfields are defined by a simple pair of first order superconstraints (2.1), while their non-minimal brethren are defined by one simple second order superconstraint (2.19). Quartoid superfields are defined by a simple triplet of first order superconstraints (2.3). All quartoid superfields (2.3), and two of six ‘minimal’ haploid superfields, those defined in (2.1e, f), are unidexterous (depend only on one world-sheet light-cone coordinate). Numerous other types of constrained superfields are discussed, albeit not as systematically, in § 2.6.

The possible choices for a Lagrangian density for the haploid and quartoid superfields are discussed in § 3. The otherwise infinite number of choices is curbed by demanding that the resulting equations of motion contain no higher than second derivatives on bosons, and no higher than first derivatives on fermions, and that no coefficient function turns out to have negative canonical dimension. This leads to the general form of the Lagrangian density (3.4), with terms supplied in (3.5), (3.9), (3.18), (3.22) and (3.24). The exhaustive construction and verification of supersymmetry of these Lagrangian density terms was facilitated by creating lists of (superderivative) superfields which are annihilated by

some subset of the four superderivatives (2.29) and (2.30). §4 discusses some issues of quantization of constrained superfields.

Upon choosing a preferred Lagrangian density, one can turn to a study of the Hilbert space, the structure of the supersymmetric wave-functions, correlation functions, and the relations to the geometry of the target space. This is illustrated in §5, where some of the basic results are recalled and generalized.

6.1. Further topics

En route, a number of issues have been deferred to a later study, and we recall some of these, for the sake of the Reader.

Transformations and ‘dualities’

In §1.3, the group of discrete transformations of the ‘soul’ of the (2,2)-superspace (generated by the four fermionic coordinates) has been studied. Two of these transformations have been identified as the root of two of the dualities in string theory:

1. left-handed conjugation, $\mathbf{C}_+$, as being the root of mirror symmetry, and
2. unconjugate parity, $\mathbf{p}$, as being the root of the Type IIA $\leftrightarrow$ Type IIB duality [15].

Rather conspicuously, mirror symmetry is an actual symmetry of the total Hilbert space, while the Type IIA $\leftrightarrow$ Type IIB duality is not. At the same time, $\mathbf{C}_+$ commutes with Lorentz symmetry while $\mathbf{p}$ does not; more importantly, $\mathbf{C}_+$ is a symmetry of the supersymmetry algebra (1.6), while $\mathbf{p}$ is not. It is tempting to conjecture that this latter property of $\mathbf{p}$ makes it the root of a duality between two *different* string theories, rather than a symmetry of a single (super)string theory, as is the case with $\mathbf{C}_+$ and mirror symmetry. In any case, these operations are but two elements of a discrete group, denoted in §1.3 as $2S_4$. The remaining 45 non-trivial operations may similarly generate additional ‘duality’ (‘triality’, and even ‘quadriality’) relations among (super)string (and related) theories, and are well worth exploring. In particular, S_4 is generated by one more independent simple swap. This third simple transformation (involution, *i.e.*, self-inverse) should at least be of interest in (super)string theory. As noted, there is however a continuous group of transformations, $GL(4, \mathbb{C})$, at least a subgroup of which could provide further insight into (super)string (and related) dynamics. This too seems to warrant a closer study.

Dimensional reduction

Supersymmetric quantum mechanics provides the underlying framework for understanding the general features of any supersymmetric field theory [19]. The present analysis of (2,2)-supersymmetric models in 2-dimensional spacetime can of course be dimensionally reduced to (2,2)-supersymmetric quantum mechanics. Since the Lorentz symmetry in 2-dimensional spacetimes is already abelian, it is not clear if any generality would be lost in this way. That is, it remains unclear whether a similarly intrinsic (*ab initio*) and comprehensive study of supersymmetric quantum mechanics would allow for possibilities not obtained by dimensional reduction from the results presented and indicated herein. The study in Refs. [28] does seem to suggest this, in fact, $\aleph_0$ -fold so.

On the other hand, with no continuous Lorentz symmetry in supersymmetric quantum mechanics, the notion of chirality is now lost. Then, the spin labels ($\pm$ sub- and super-scripts) as used in this article simply count the extended supersymmetries. This change has to be taken into account when translating the present results to various applications of $N \geq 4$ supersymmetric quantum mechanics.

Quantization

Some of the issues of quantization of constrained superfields have been mentioned in § 4. A complete and detailed Hamiltonian (Dirac, BRST, *etc.*) treatment would seem to be well worth, especially in view of the apparent difficulties with unidexterous (super)fields; see above, and Ref. [10] for a recent account and further references. Suffice it here to note that both general approaches outlined in § 4 have both advantages and drawbacks. While the use of unconstrained superfields allows for a more straightforward quantization, it also requires many more component fields and a complementing set of gauge symmetries (to gauge away the extra components introduced by passing to unconstrained superfields).

On the other hand, quantization of constrained superfields requires a detailed study of the full content of the constraints, and a proper enforcement by means of suitable Lagrange multipliers. As the component fields of the Lagrange multiplier superfields ultimately must disappear from the action, the proper Siegel (κ -) symmetry must be identified and ensured.

Gauge symmetries

As noted from the outset, gauge symmetry issues (in a comparably comprehensive setting) are completely ignored herein, and are deferred to a later effort. Partial results may be found in Refs. [3,4].

However, quite unrelated to usual gauge symmetries, Eqs. (2.22) present a rather peculiar type of inhomogeneous ‘gauge’ transformations, where the supercoordinates themselves serve as generators.

These are perhas not unrelated to the relation:

$$\begin{aligned} \{\Phi : \bar{D}_{\pm}\Phi=0\} &\simeq \{\overset{\circ}{\Phi} \simeq \overset{\circ}{\Phi} + \bar{\varsigma}^+ \overset{\circ}{\Phi}_+^{(1)} + \bar{\varsigma}^- \overset{\circ}{\Phi}_-^{(1)}\} , \\ \text{where } \overset{\circ}{\Phi}_+^{(1)} &\simeq \overset{\circ}{\Phi}_+^{(1)} + \bar{\varsigma}^- \overset{\circ}{\Phi}^{(2)} , \quad \overset{\circ}{\Phi}_-^{(1)} \simeq \overset{\circ}{\Phi}_-^{(1)} + \bar{\varsigma}^+ \overset{\circ}{\Phi}^{(2)} . \end{aligned} \tag{6.1}$$

Here, as throughout the article, the circle atop indicates that the superfield is unconstrained. The supefields $\Phi^{(i)}$ may be regarded as i^{th} order gauge parameters. That is, component fields of the originally unconstrained superfield $\overset{\circ}{\Phi}$ are being eliminated by means of the (1st order) gauge transformation

$$\delta \overset{\circ}{\Phi} = \bar{\varsigma}^+ \overset{\circ}{\Phi}_+^{(1)} + \bar{\varsigma}^- \overset{\circ}{\Phi}_-^{(1)} . \tag{6.2a}$$

This, however, would attempt to gauge away some of the component fields twice (recall the doubly shaded region in Figs. 4 and 6), and so the gauge parameters, $\overset{\circ}{\Phi}_{\pm}^{(1)}$, are themselves subject to a (2nd order) gauge transformation,

$$\delta \overset{\circ}{\Phi}_-^{(1)} = \bar{\varsigma}^+ \overset{\circ}{\Phi}^{(2)} , \quad \text{and} \quad \delta \overset{\circ}{\Phi}_+^{(1)} = \bar{\varsigma}^- \overset{\circ}{\Phi}^{(2)} , \tag{6.2b}$$

with respect to which the first order gauge transformation (6.2a) is invariant. The component fields remaining in this sequential double quotient are the same as those in a chiral superfield. This sequential double quotient representation of the chiral superfield is obviously rather more involved technically. However, it also gives the chiral superfield the rôle of a gauge field (having an inhomogeneous transformation), which is certainly not the usual case. The non-minimal chiral superfield also admits a similar description:

$$\{\Theta : \bar{D}_+ \bar{D}_- \Theta = 0\} \simeq \{\mathring{\Theta} \simeq \mathring{\Theta} + \bar{\varsigma}^- \bar{\varsigma}^+ \mathring{\Theta}^{(1)}\} , \quad (6.3)$$

which is a simple quotient by the inhomogeneous transformation

$$\delta \mathring{\Theta} = \bar{\varsigma}^- \bar{\varsigma}^+ \mathring{\Theta}^{(1)} , \quad (6.4)$$

as compared to the sequential double quotient, with respect to the cascading pair of inhomogeneous transformations (6.2), of the ‘minimal’ chiral superfield. Of course, the other haploid superfields, ‘minimal’ and non-minimal, admit similar quotient representations. Quite generally, the constrained and the quotient representations are ‘dual’ in a well-defined sense: the former represents the kernel, the latter the cokernel of a mapping.

The utility of this quotient representation as well as the duality between the quotient and the constrained representations remains unclear, and will require a study in its own right. It is however clear that it will be linked to the possible rôles of the ‘peculiar’ inhomogeneous transformations, such as (2.22), (6.2) and (6.4). Note also that the quotient description has a direct implication in a BRST description: Eq. (6.4) induces a single set of ghost superfields, whereas Eqs. (6.2a, b) induce a set of ghost superfields, and a set of ghost-for-ghost superfields, respectively.

Other superconstraints

Finally, as shown in § 2.6, the simple superconstraints (2.1), (2.3) and (2.19) barely scratch the surface of the incredibly rich category of superconstraint theory. While the abelian nature of Lorentz symmetry in ≤ 2 -dimensional spacetime is the major reason of the much larger number of possibilities for defining superconstraints, several of the truly novel superconstraints of § 2.6 are equally well possible in > 2 -dimensional spacetimes! Unfortunately, no natural or systematic way of enlisting and studying these non-simple, non-homogeneous or non-linear superconstraints seems to present itself, and it is not even clear that any *finite* form of enlisting is even possible or whether such efforts are worth the trouble.

Clearly, however, some of these do deserve a closer study, such as the family of superconstraints (2.49), which interpolates continuously between chiral and twisted-chiral superfields, and hence between models the IR regime of which is believed to correspond to mirror compactifications in string theory. A similar family interpolating between $\mathbf{p}$ -reflected models does not seem to be possible in 2-dimensions, as $\mathbf{p}$ does not commute with Lorentz symmetry. Even in supersymmetric quantum mechanics, a construction of such an interpolation is obstructed because $\mathbf{p}$ does not leave the superalgebra (1.6) invariant. This seems to further deepen the chasm between mirror symmetry, as generated by $\mathbf{C}_+$, and the Type IIA $\leftrightarrow$ Type IIB duality, as generated by $\mathbf{p}$. These and related issues are left for a subsequent study.

Appendix A. Notational Nitpicking

Following Wess and Bagger's definitions [6]:

$$\psi^\alpha \stackrel{\text{def}}{=} \epsilon^{\alpha\beta} \psi_\beta, \quad \bar{\psi}^{\dot{\alpha}} \stackrel{\text{def}}{=} \epsilon^{\dot{\alpha}\dot{\beta}} \bar{\psi}_{\dot{\beta}}, \quad \psi_\alpha = \epsilon_{\alpha\beta} \psi^\beta, \quad \bar{\psi}_{\dot{\alpha}} = \epsilon_{\dot{\alpha}\dot{\beta}} \bar{\psi}^{\dot{\beta}}, \quad (\text{A.1})$$

and

$$\psi \cdot \chi \stackrel{\text{def}}{=} \psi^\alpha \chi_\alpha, \quad \text{but} \quad \bar{\psi} \cdot \bar{\chi} \stackrel{\text{def}}{=} \bar{\psi}_{\dot{\alpha}} \bar{\chi}^{\dot{\alpha}}. \quad (\text{A.2})$$

It follows that

$$\varsigma^2 \stackrel{\text{def}}{=} +\epsilon_{\alpha\beta} \varsigma^\alpha \varsigma^\beta = 2\varsigma^+ \varsigma^-, \quad D^2 \stackrel{\text{def}}{=} \epsilon^{\alpha\beta} D_\beta D_\alpha = 2D_+ D_- , \quad (\text{A.3a})$$

but

$$\bar{\varsigma}^2 \stackrel{\text{def}}{=} +\epsilon_{\dot{\alpha}\dot{\beta}} \bar{\varsigma}^{\dot{\alpha}} \bar{\varsigma}^{\dot{\beta}} = 2\bar{\varsigma}^- \bar{\varsigma}^+, \quad \bar{D}^2 \stackrel{\text{def}}{=} \epsilon^{\dot{\alpha}\dot{\beta}} \bar{D}_{\dot{\beta}} \bar{D}_{\dot{\alpha}} = 2\bar{D}_- \bar{D}_+ . \quad (\text{A.3b})$$

Therefore (recall Eqs. (1.5)):

$$D_\alpha \varsigma^\beta = \delta_\alpha^\beta, \quad D^2 \varsigma^2 = -4, \quad \bar{D}_{\dot{\alpha}} \bar{\varsigma}^{\dot{\beta}} = -\delta_{\dot{\alpha}}^{\dot{\beta}}, \quad \bar{D}^2 \bar{\varsigma}^2 = -4. \quad (\text{A.4})$$

Notice that Eqs. (A.1) imply that

$$\psi^- = \psi_+, \quad \psi^+ = -\psi_-, \quad \bar{\psi}^- = \bar{\psi}_+, \quad \bar{\psi}^+ = -\bar{\psi}_-, \quad (\text{A.5})$$

We say that $\psi^- = \psi_+$ (and their conjugates) have spin $-\frac{1}{2}$, and $\psi^+ = -\psi_-$ (and their conjugates) have spin $+\frac{1}{2}$.

Fermionic Dirac delta-functions are utterly simple: *e.g.*, $\delta(\varsigma^+) \equiv \varsigma^+$. For multiple delta-functions, we must merely choose an order. Fixing $\delta^4(\varsigma) \stackrel{\text{def}}{=} \bar{\varsigma}^- \bar{\varsigma}^+ \varsigma^+ \varsigma^-$ so that $\int d^4\varsigma \delta^4(\varsigma) = 1$, we find:

$$\delta^2(\varsigma) = \varsigma^+ \varsigma^-, \quad \delta^2(\not{\varsigma}) = \bar{\varsigma}^+ \varsigma^-, \quad \delta^2(\varsigma^\pm) = \bar{\varsigma}^- \varsigma^\pm, \quad (\text{A.6})$$

$$\delta^2(\bar{\varsigma}) = \bar{\varsigma}^- \bar{\varsigma}^+, \quad \delta^2(\not{\bar{\varsigma}}) = \bar{\varsigma}^- \bar{\varsigma}^+, \quad \delta^2(\bar{\varsigma}^\pm) = \bar{\varsigma}^\pm \bar{\varsigma}^+. \quad (\text{A.7})$$

The following *operatorial* identities may be of help:

$$[D_+, \bar{D}_+] \equiv 2(D_+ \bar{D}_+ - i\partial_+), \quad [D_-, \bar{D}_-] \equiv 2(D_- \bar{D}_- - i\partial_-), \quad (\text{A.8})$$

whence

$$\begin{aligned} [D_-, \bar{D}_-][D_+, \bar{D}_+] &\equiv 2[D_-, \bar{D}_-](D_+ \bar{D}_+ - i\partial_+) , \\ &\equiv 2[D_-, \bar{D}_-]D_+ \bar{D}_+ - 4i\partial_+ D_- \bar{D}_- - 4\partial_+ \partial_- , \\ &\equiv 2[D_+, \bar{D}_+](D_- \bar{D}_- - i\partial_-) , \\ &\equiv 2[D_+, \bar{D}_+]D_- \bar{D}_- - 4i\partial_- D_+ \bar{D}_+ - 4\partial_+ \partial_- . \end{aligned} \quad (\text{A.9})$$

Also,

$$\begin{aligned} [D_-, \bar{D}_-]D_- &\equiv 2i\partial_- D_- , & [D_-, \bar{D}_-]\bar{D}_- &\equiv -2i\partial_- \bar{D}_- , \\ [D_+, \bar{D}_+]D_+ &\equiv 2i\partial_+ D_+ , & [D_+, \bar{D}_+]\bar{D}_+ &\equiv -2i\partial_+ \bar{D}_+ , \end{aligned} \quad (\text{A.10})$$

and

$$([D_1, D_2]AB) = ([D_1, D_2]A)B + A([D_1, D_2]B) + (-)^A 2(D_{[1}A)(D_{2]}B), \quad (\text{A.11})$$

for any two superderivatives D_1, D_2 and any two superfields A, B , and where

$$(-)^A \stackrel{\text{def}}{=} \begin{cases} +1 & \text{if } A \text{ is commuting,} \\ -1 & \text{anticommuting.} \end{cases} \quad (\text{A.12})$$

Appendix B. Expounding Expansions

For the benefit of the Reader unskilled in projecting onto component fields by means of the superderivatives in § 2.3, basic definitions and some simple consequences are listed here.

B.1. Component fields

Chiral superfield: $\bar{D}_\pm \Phi = 0$

$$\begin{aligned} \Phi| &= \phi, \quad D_\pm \Phi| = \sqrt{2}\psi_\pm, \quad D_- D_+ \Phi| = 2F, \\ [D_-, \bar{D}_-] \Phi| &= -\bar{D}_- D_- \Phi| = -2i(\partial_- \phi), \\ [D_+, \bar{D}_+] \Phi| &= -\bar{D}_+ D_+ \Phi| = -2i(\partial_+ \phi), \\ [D_-, \bar{D}_-] D_+ \Phi| &= -\bar{D}_- D_- D_+ \Phi| = -2i(\partial_- \psi_+), \\ [D_+, \bar{D}_+] D_- \Phi| &= -\bar{D}_+ D_+ D_- \Phi| = -2i(\partial_+ \psi_-), \\ [D_-, \bar{D}_-][D_+, \bar{D}_+] \Phi| &= -[D_-, \bar{D}_-] \bar{D}_+ D_+ \Phi| = -[D_+, \bar{D}_+] \bar{D}_- D_- \Phi| = -4(\partial_+ \partial_- \phi). \end{aligned}$$

Antichiral superfield: $D_\pm \bar{\Phi} = 0$

$$\begin{aligned} \bar{\Phi}| &= \phi, \quad \bar{D}_\pm \bar{\Phi}| = \sqrt{2}\bar{\psi}_\pm, \quad \bar{D}_+ \bar{D}_- \bar{\Phi}| = 2\bar{F}, \\ [D_-, \bar{D}_-] \bar{\Phi}| &= D_- \bar{D}_- \bar{\Phi}| = 2i(\partial_- \bar{\phi}), \\ [D_+, \bar{D}_+] \bar{\Phi}| &= D_+ \bar{D}_+ \bar{\Phi}| = 2i(\partial_+ \bar{\phi}), \\ [D_-, \bar{D}_-] \bar{D}_+ \bar{\Phi}| &= D_- \bar{D}_- \bar{D}_+ \bar{\Phi}| = 2i(\partial_- \bar{\psi}_+), \\ [D_+, \bar{D}_+] \bar{D}_- \bar{\Phi}| &= D_+ \bar{D}_+ \bar{D}_- \bar{\Phi}| = 2i(\partial_+ \bar{\psi}_-), \\ [D_-, \bar{D}_-][D_+, \bar{D}_+] \bar{\Phi}| &= [D_-, \bar{D}_-] D_+ \bar{D}_+ \bar{\Phi}| = [D_+, \bar{D}_+] D_- \bar{D}_- \bar{\Phi}| = -4(\partial_+ \partial_- \bar{\phi}). \end{aligned}$$

Twisted-chiral superfield: $\bar{D}_- \Xi = 0 = D_+ \Xi$

$$\begin{aligned} \Xi| &= x, \quad D_- \Xi| = \sqrt{2}\xi_-, \quad \bar{D}_+ \Xi| = \sqrt{2}\chi_+, \quad D_- \bar{D}_+ \Xi| = 2X, \\ [D_-, \bar{D}_-] \Xi| &= -\bar{D}_- D_- \Xi| = -2i(\partial_- x), \\ [D_+, \bar{D}_+] \Xi| &= D_+ \bar{D}_+ \Xi| = 2i(\partial_+ x), \\ [D_-, \bar{D}_-] \bar{D}_+ \Xi| &= -\bar{D}_- D_- \bar{D}_+ \Xi| = -2i(\partial_- \chi_+), \\ [D_+, \bar{D}_+] D_- \Xi| &= D_+ \bar{D}_+ D_- \Xi| = 2i(\partial_+ \xi_-), \\ [D_-, \bar{D}_-][D_+, \bar{D}_+] \Xi| &= [D_-, \bar{D}_-] D_+ \bar{D}_+ \Xi| = -[D_+, \bar{D}_+] \bar{D}_- D_- \Xi| = 4(\partial_+ \partial_- \phi). \end{aligned}$$

Twisted-antichiral superfield: $D_- \bar{\Xi} = 0 = \bar{D}_+ \bar{\Xi}$

$$\begin{aligned} \bar{\Xi}| &= \phi, \quad \bar{D}_- \bar{\Xi}| = \sqrt{2}\bar{\xi}_-, \quad D_+ \bar{\Xi}| = \sqrt{2}\bar{\chi}_+, \quad D_+ \bar{D}_- \bar{\Xi}| = 2\bar{X}, \\ [D_-, \bar{D}_-] \bar{\Xi}| &= D_- \bar{D}_- \bar{\Xi}| = 2i(\partial_- \bar{x}), \\ [D_+, \bar{D}_+] \bar{\Xi}| &= -\bar{D}_+ D_+ \bar{\Xi}| = -2i(\partial_+ \bar{x}), \\ [D_-, \bar{D}_-] D_+ \bar{\Xi}| &= D_- \bar{D}_- D_+ \bar{\Xi}| = 2i(\partial_- \bar{\chi}_+), \end{aligned}$$

$$\begin{aligned}
[D_+, \bar{D}_+] \bar{D}_- \Xi &= -\bar{D}_+ D_+ \bar{D}_- \Xi = -2i(\partial_{\pm} \bar{\xi}_-), \\
[D_-, \bar{D}_-][D_+, \bar{D}_+] \Xi &= -[D_-, \bar{D}_-] \bar{D}_+ D_+ \Xi = [D_+, \bar{D}_+] D_- \bar{D}_- \Xi = 4(\partial_{\pm} \partial_{\pm} \bar{x}).
\end{aligned}$$

Lefton superfield: $D_- \Lambda = 0 = \bar{D}_- \Lambda$

$$\begin{aligned}
\Lambda &= \ell, \quad D_+ \Lambda = \sqrt{2} \lambda_+, \quad \bar{D}_+ \Lambda = \sqrt{2} \chi_+, \\
[D_+, \bar{D}_+] \Lambda &= 4L_{\pm}, \quad D_+ \bar{D}_+ \Lambda = i(\partial_{\pm} \ell) + 2L_{\pm}, \quad \bar{D}_+ D_+ \Lambda = i(\partial_{\pm} \ell) - 2L_{\pm}.
\end{aligned}$$

Righton superfield: $D_+ \Upsilon = 0 = \bar{D}_+ \Upsilon$

$$\begin{aligned}
\Upsilon &= x, \quad D_- \Upsilon = \sqrt{2} \xi_-, \quad \bar{D}_+ \Upsilon = \sqrt{2} \chi_+, \\
[D_-, \bar{D}_-] \Upsilon &= 4R_{\pm}, \quad D_- \bar{D}_- \Upsilon = i(\partial_{\pm} r) + 2R_{\pm}, \quad \bar{D}_- D_- \Upsilon = i(\partial_{\pm} r) - 2R_{\pm}.
\end{aligned}$$

Non-minimal chiral superfield: $\bar{D}_+ \bar{D}_- \Theta = 0$

$$\begin{aligned}
\Theta &= t, \quad D_{\pm} \Theta = \sqrt{2} \theta_{\pm}, \quad D_- D_+ \Theta = 2T, \\
[D_-, \bar{D}_-] \Theta &= 4T_{\pm}, \quad D_- \bar{D}_- \Theta = i(\partial_{\pm} t) + 2T_{\pm}, \quad D_- \bar{D}_+ \Theta = 2T_{\mp}, \\
[D_+, \bar{D}_+] \Theta &= 4T_{\pm}, \quad D_+ \bar{D}_+ \Theta = i(\partial_{\pm} t) + 2T_{\pm}, \quad D_+ \bar{D}_- \Theta = 2T_{\pm}, \\
[D_-, \bar{D}_-] D_+ \Theta &= 4\sqrt{2} \tau_-, \quad [D_-, \bar{D}_-] \bar{D}_+ \Theta = -2\sqrt{2} i(\partial_{\pm} \vartheta_+), \\
[D_+, \bar{D}_+] D_- \Theta &= 4\sqrt{2} \tau_+, \quad [D_+, \bar{D}_+] \bar{D}_- \Theta = -2\sqrt{2} i(\partial_{\pm} \vartheta_-), \\
[D_-, \bar{D}_-][D_+, \bar{D}_+] \Theta &= 4[(\partial_{\pm} \partial_{\pm} t) - 2i(\partial_{\pm} T_{\pm}) - 2i(\partial_{\pm} T_{\pm})].
\end{aligned}$$

Non-minimal antichiral superfield: $D_- D_+ \bar{\Theta} = 0$

$$\begin{aligned}
\bar{\Theta} &= \bar{t}, \quad D_{\pm} \bar{\Theta} = \sqrt{2} \bar{\theta}_{\pm}, \quad \bar{D}_+ \bar{D}_- \bar{\Theta} = 2\bar{T}, \\
[D_-, \bar{D}_-] \bar{\Theta} &= 4\bar{T}_{\pm}, \quad D_- \bar{D}_- \bar{\Theta} = i(\partial_{\pm} \bar{t}) + 2\bar{T}_{\pm}, \quad D_- \bar{D}_+ \bar{\Theta} = 2\bar{T}_{\mp}, \\
[D_+, \bar{D}_+] \bar{\Theta} &= 4\bar{T}_{\pm}, \quad D_+ \bar{D}_+ \bar{\Theta} = i(\partial_{\pm} \bar{t}) + 2\bar{T}_{\pm}, \quad D_+ \bar{D}_- \bar{\Theta} = 2\bar{T}_{\pm}, \\
[D_-, \bar{D}_-] \bar{D}_+ \bar{\Theta} &= 4\sqrt{2} \bar{\tau}_-, \quad [D_-, \bar{D}_-] D_+ \bar{\Theta} = 2\sqrt{2} i(\partial_{\pm} \bar{\vartheta}_+), \\
[D_+, \bar{D}_+] \bar{D}_- \bar{\Theta} &= 4\sqrt{2} \bar{\tau}_+, \quad [D_+, \bar{D}_+] D_- \bar{\Theta} = 2\sqrt{2} i(\partial_{\pm} \bar{\vartheta}_-), \\
[D_-, \bar{D}_-][D_+, \bar{D}_+] \bar{\Theta} &= 4[(\partial_{\pm} \partial_{\pm} \bar{t}) + 2i(\partial_{\pm} \bar{T}_{\pm}) + 2i(\partial_{\pm} \bar{T}_{\pm})].
\end{aligned}$$

Non-minimal twisted-chiral superfield: $D_+ \bar{D}_- \Pi = 0$

$$\begin{aligned}
\Pi &= p, \quad D_{\pm} \Pi = \sqrt{2} \pi_{\pm}, \quad \bar{D}_{\pm} \Pi = \sqrt{2} \varpi_{\pm}, \quad D_- D_+ \Pi = 2P, \\
[D_-, \bar{D}_-] \Pi &= 4P_{\pm}, \quad D_- \bar{D}_- \Pi = i(\partial_{\pm} p) + 2P_{\pm}, \quad D_- \bar{D}_+ \Pi = 2P_{\mp}, \\
[D_+, \bar{D}_+] \Pi &= 4P_{\pm}, \quad D_+ \bar{D}_+ \Pi = i(\partial_{\pm} p) + 2P_{\pm}, \quad \bar{D}_+ \bar{D}_- \Pi = 2P, \\
[D_-, \bar{D}_-] D_+ \Pi &= -2\sqrt{2} i(\partial_{\pm} \pi_+), \quad [D_-, \bar{D}_-] \bar{D}_+ \Pi = 4\sqrt{2} \tilde{\varphi}_-, \\
[D_+, \bar{D}_+] D_- \Pi &= 4\sqrt{2} \varphi_+, \quad [D_+, \bar{D}_+] \bar{D}_- \Pi = 2\sqrt{2} i(\partial_{\pm} \varpi_-), \\
[D_-, \bar{D}_-][D_+, \bar{D}_+] \Pi &= -4[(\partial_{\pm} \partial_{\pm} p) - 2i(\partial_{\pm} P_{\pm}) + 2i(\partial_{\pm} P_{\pm})].
\end{aligned}$$

Non-minimal twisted-antichiral superfield: $D_- \bar{D}_+ \bar{\Pi} = 0$

$$\begin{aligned}
\bar{\Pi} &= \bar{p}, \quad D_{\pm} \bar{\Pi} = \sqrt{2} \bar{\pi}_{\pm}, \quad \bar{D}_{\pm} \bar{\Pi} = \sqrt{2} \bar{\varpi}_{\pm}, \quad D_- D_+ \bar{\Pi} = 2\bar{P}, \\
[D_-, \bar{D}_-] \bar{\Pi} &= 4\bar{P}_{\pm}, \quad D_- \bar{D}_- \bar{\Pi} = i(\partial_{\pm} \bar{p}) + 2\bar{P}_{\pm}, \quad D_- \bar{D}_+ \bar{\Pi} = 2\bar{P}_{\mp}, \\
[D_+, \bar{D}_+] \bar{\Pi} &= 4\bar{P}_{\pm}, \quad D_+ \bar{D}_+ \bar{\Pi} = i(\partial_{\pm} \bar{p}) + 2\bar{P}_{\pm}, \quad \bar{D}_+ \bar{D}_- \bar{\Pi} = 2\bar{P}, \\
[D_-, \bar{D}_-] D_+ \bar{\Pi} &= 4\sqrt{2} \tilde{\varphi}_-, \quad [D_-, \bar{D}_-] \bar{D}_+ \bar{\Pi} = 2\sqrt{2} i(\partial_{\pm} \bar{\pi}_+), \\
[D_+, \bar{D}_+] D_- \bar{\Pi} &= -2\sqrt{2} i(\partial_{\pm} \bar{\varpi}_-), \quad [D_+, \bar{D}_+] \bar{D}_- \bar{\Pi} = 4\sqrt{2} \bar{\varphi}_+, \\
[D_-, \bar{D}_-][D_+, \bar{D}_+] \bar{\Pi} &= -4[(\partial_{\pm} \partial_{\pm} \bar{p}) - 2i(\partial_{\pm} \bar{P}_{\pm}) + 2i(\partial_{\pm} \bar{P}_{\pm})].
\end{aligned}$$

B.2. Projection technique

As an example of the projection technique, below is the detailed expansion of the λ -term $\frac{1}{8}[D_+, \bar{D}_+]D_-(\bar{D}_-\Phi^{\bar{\nu}})M_{\bar{\nu}}$, which has been discussed in §3.4. The uninitiated Reader should be able to follow each step and then apply this technique for any of the other parts of the Lagrangian density (3.4).

$$\begin{aligned} & \frac{1}{8}[D_+, \bar{D}_+]D_-(\bar{D}_-\Phi^{\bar{\nu}})M_{\bar{\nu}} \\ &= \frac{1}{8}[D_+, \bar{D}_+][(2i\partial_=\Phi^{\bar{\nu}})M_{\bar{\nu}} + (D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})M_{\bar{\nu},\mu}] , \end{aligned} \quad (\text{B.1a})$$

$$\begin{aligned} &= \frac{1}{8}\left\{ (2i\partial_=[D_+, \bar{D}_+]\Phi^{\bar{\nu}})M_{\bar{\nu}} + (2i\partial_=\Phi^{\bar{\nu}})([D_+, \bar{D}_+]M_{\bar{\nu}}) \right. \\ &\quad + 2(2i\partial_=\underline{D_+\Phi^{\bar{\nu}}})(\bar{D}_+M_{\bar{\nu}}) - 2(2i\partial_=\bar{D}_+\Phi^{\bar{\nu}})(D_+M_{\bar{\nu}}) \\ &\quad + [[D_+, \bar{D}_+](D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})]M_{\bar{\nu},\mu} + (D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})([D_+, \bar{D}_+]M_{\bar{\nu},\mu}) \\ &\quad \left. + 2[D_+(D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})](\bar{D}_+M_{\bar{\nu},\mu}) - 2[\bar{D}_+(D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})](D_+M_{\bar{\nu},\mu}) \right\} \quad (\text{B.1b}) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{8}\left\{ (2i\partial_=(2i\partial_{\mp}\Phi^{\bar{\nu}}))M_{\bar{\nu}} \right. \\ &\quad + (2i\partial_=\Phi^{\bar{\nu}})[D_+\bar{D}_+(\bar{\Phi}_L + \Xi) - (\bar{D}_+D_+\Phi) + [D_+, \bar{D}_+]\Lambda]M'_{\bar{\nu}} \\ &\quad \quad + [D_+(\Phi + \Lambda), \bar{D}_+(\bar{\Phi}_L + \Xi + \Lambda)]M''_{\bar{\nu}} \\ &\quad - 4i(\partial_=\bar{D}_+\Phi^{\bar{\nu}})[D_+(\Phi + \Lambda)]M'_{\bar{\nu}} \\ &\quad + (([D_+, \bar{D}_+]D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}}) + (D_-\Phi^{\mu})([D_+, \bar{D}_+]\bar{D}_-\Phi^{\bar{\nu}}) \\ &\quad \quad - 2(D_+D_-\Phi^{\mu})(\bar{D}_+\bar{D}_-\Phi^{\bar{\nu}}) + 2(\underline{\bar{D}_+D_-\Phi^{\mu}})(\underline{D_+\bar{D}_-\Phi^{\bar{\nu}}}))M_{\bar{\nu},\mu} \\ &\quad + (D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}})[D_+\bar{D}_+(\bar{\Phi}_L + \Xi) - (\bar{D}_+D_+\Phi) + [D_+, \bar{D}_+]\Lambda]M'_{\bar{\nu},\mu} \\ &\quad \quad + [D_+(\Phi + \Lambda), \bar{D}_+(\bar{\Phi}_L + \Xi + \Lambda)]M''_{\bar{\nu},\mu} \\ &\quad + 2[(D_+D_-\Phi^{\mu})(\bar{D}_-\Phi^{\bar{\nu}}) - (D_-\Phi^{\mu})(\underline{D_+\bar{D}_-\Phi^{\bar{\nu}}})][\bar{D}_+(\bar{\Phi}_L + \Xi + \Lambda)]M'_{\bar{\nu},\mu} \\ &\quad \left. - 2[(\underline{\bar{D}_+D_-\Phi^{\mu}})(\bar{D}_-\Phi^{\bar{\nu}}) - (D_-\Phi^{\mu})(\bar{D}_+\bar{D}_-\Phi^{\bar{\nu}})][D_+(\Phi + \Lambda)]M'_{\bar{\nu},\mu} \right\} , \quad (\text{B.1c}) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{8}\left\{ -4(\partial_=\partial_{\mp}\phi^{\bar{\nu}})M_{\bar{\nu}} \right. \\ &\quad + (2i\partial_=\phi^{\bar{\nu}})[[2i\partial_{\mp}(\bar{\phi}_L + x - \phi) + 4L_{\mp}]M'_{\bar{\nu}} + 4(\psi_+ + \lambda_+)(\bar{\psi}_L + \chi_+ + \lambda_+)M''_{\bar{\nu}}] \\ &\quad - 8i(\partial_=\psi_+^{\bar{\nu}})(\psi_+ + \lambda_+)M'_{\bar{\nu}} \\ &\quad + [2(-2i\partial_{\mp}\psi_-^{\mu})(\psi_-^{\bar{\nu}}) + 2(\psi_-^{\mu})(2i\partial_{\mp}\psi_-^{\bar{\nu}}) - 2(-2F^{\mu})(2F^{\bar{\nu}})]M_{\bar{\nu},\mu} \\ &\quad + 2\psi_-^{\mu}\psi_-^{\bar{\nu}}[[2i\partial_{\mp}(\bar{\phi}_L + x - \phi) + 4L_{\mp}]M'_{\bar{\nu},\mu} + 4(\psi_+ + \lambda_+)(\bar{\psi}_L + \chi_+ + \lambda_+)M''_{\bar{\nu},\mu}] \\ &\quad \left. + 4(-2F^{\mu})\psi_-^{\bar{\nu}}(\bar{\psi}_L + \chi_+ + \lambda_+)M'_{\bar{\nu},\mu} + 8\psi_-^{\mu}(F^{\bar{\nu}})(\psi_+ + \lambda_+)M'_{\bar{\nu},\mu} \right\} , \quad (\text{B.1d}) \end{aligned}$$

We have immediately used that $(D_-\bar{D}_-\bar{\Phi}) = (2i\partial_=\bar{\Phi})$, and underlined the terms that

vanish because of the definitions $\bar{D}_\pm \Phi = 0$ and $D_\pm \bar{\Phi} = 0$. Terms such as $[D_+(\Phi + \Lambda)]M'_\nu$ are simply an abbreviation for

$$[D_+(\Phi + \Lambda)]M'_\nu \rightarrow [(D_+\Phi^\rho)M_{\bar{\nu},\rho} + (D_+\Lambda^{\hat{a}})M_{\bar{\nu},\hat{a}}] . \quad (\text{B.2})$$

Next, use that

$$-4(\partial_=\partial_\mp\phi^{\bar{\nu}})M_{\bar{\nu}} = -4\partial_\mp[(\partial_=\phi^{\bar{\nu}})M_{\bar{\nu}}] + 4(\partial_=\phi^{\bar{\nu}})[\partial_\mp(\bar{\phi}_L+x+\phi)]M'_\nu , \quad (\text{B.3})$$

and drop a few total derivative terms to obtain (after some straightforward algebra):

$$\begin{aligned} & \frac{1}{8}[D_+, \bar{D}_+]D_-(\bar{D}_-\Phi^{\bar{\nu}})M_{\bar{\nu}}| \\ & \cong [(\partial_=\phi^{\bar{\nu}})(\partial_\mp\phi^\mu) + F^\mu F^{\bar{\nu}} + \frac{i}{2}(\psi_+^{\bar{\nu}}\overleftrightarrow{\partial}_=\psi_+^\mu) + \frac{i}{2}(\psi_-^{\bar{\nu}}\overleftrightarrow{\partial}_\mp\psi_-^\mu)]M_{\bar{\nu},\mu} \end{aligned} \quad (\text{B.4a})$$

$$+ [(\partial_=\phi^{\bar{\nu}})[\frac{1}{2}(\partial_\mp\ell^{\hat{a}}) + iL_\mp] + \frac{i}{2}(\psi_+^{\bar{\nu}}\overleftrightarrow{\partial}_=\lambda_+^{\hat{a}})]M_{\bar{\nu},\hat{a}} \quad (\text{B.4b})$$

$$+ i(\partial_=\phi^{\bar{\nu}})[\psi_+^\rho(\bar{\psi}_{L+}^{\tilde{\mu}}M_{\bar{\nu},\tilde{\mu}\rho} + \chi_+^\alpha M_{\bar{\nu},\alpha\rho} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\rho}) \quad (\text{B.4c})$$

$$+ \lambda_+^{\hat{b}}(\bar{\psi}_{L+}^{\tilde{\mu}}M_{\bar{\nu},\tilde{\mu}\hat{b}} + \chi_+^\alpha M_{\bar{\nu},\alpha\hat{b}} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\hat{b}})] \quad (\text{B.4d})$$

$$+ \frac{i}{2}\psi_-^\rho\psi_-^{\bar{\nu}}[(\partial_\mp\bar{\phi}_L^{\tilde{\mu}})M_{\bar{\nu},\tilde{\mu}\rho} + (\partial_\mp x^\alpha)M_{\bar{\nu},\alpha\rho} - (\partial_\mp\phi^\mu)M_{\bar{\nu},\mu\rho}] + \psi_-^\rho\psi_-^{\bar{\nu}}L_\mp^{\hat{a}}M_{\bar{\nu},\hat{a}\rho} \quad (\text{B.4e})$$

$$- F^\rho\psi_-^{\bar{\nu}}(\bar{\psi}_{L+}^{\tilde{\mu}}M_{\bar{\nu},\tilde{\mu}\rho} + \chi_+^\alpha M_{\bar{\nu},\alpha\rho} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\rho}) + \psi_-^\rho F^{\bar{\nu}}(\psi_+^\mu M_{\bar{\nu},\mu\rho} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\rho}) \quad (\text{B.4f})$$

$$+ \psi_-^\sigma\psi_-^{\bar{\nu}}[\psi_+^\rho(\bar{\psi}_{L+}^{\tilde{\mu}}M_{\bar{\nu},\tilde{\mu}\rho\sigma} + \chi_+^\alpha M_{\bar{\nu},\alpha\rho\sigma} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\rho\sigma}) \quad (\text{B.4g})$$

$$+ \lambda_+^{\hat{b}}(\bar{\psi}_{L+}^{\tilde{\mu}}M_{\bar{\nu},\tilde{\mu}\hat{b}\sigma} + \chi_+^\alpha M_{\bar{\nu},\alpha\hat{b}\sigma} + \lambda_+^{\hat{a}}M_{\bar{\nu},\hat{a}\hat{b}\sigma})] . \quad (\text{B.4h})$$

Of course, the Lagrangian density must include also the hermitian conjugate terms.

Besides the contribution to the ‘standard’ D-terms, leading with the terms in (B.4a), notice the mixed kinetic terms in (B.4b), the mixed 4-fermion (Fermi or Thirring) terms in (B.4g–h), and the terms in (B.4f) which change the (still algebraic, *i.e.*, non-dynamical) equations of motion for $F, \bar{F}$. Upon ellimination of these, the supersymmetry transformation rules become non-linear in the fermionic fields, as discussed in the last part of § 3.7, and the terms in (B.4f) modify this non-linearity.

Acknowledgments: I am indebted to S.James Gates, Jr., George Minić and Joe Polchinski for helpful and encouraging discussions, and to the generous support of the Department of Energy through the grant DE-FG02-94ER-40854. Many thanks to the Institute for Theoretical Physics at Santa Barbara, where this work was completed, and the National Science Foundation for their support under Grant No. PHY94-07194.

References

- [1] S.J. Gates, Jr., C.M. Hull and M. Roček: *Nucl. Phys.* **B248** (1984)157.
- [2] E. Witten: *Nucl. Phys.* **B403** (1993)159–222.
- [3] S.J. Gates, Jr.: *Phys. Lett.* **B352** (1995)43–49.
- [4] S.J. Gates, Jr., M.T. Grisaru and M.E. Wehlau: *Nucl. Phys.* **B460** (1996)579–614.
- [5] A. Sevrin and J. Troost: *Nucl. Phys.* **B492** (1997)623–646;
M.T. Grisaru, A. Massar, A. Sevrin and J. Troost: *Fortsch. Phys.* **47**(1999)301–307.
- [6] J. Wess and J. Bagger: *Supersymmetry and Supergravity*
(Princeton University Pub., Princeton NJ, 1983).
- [7] S.J. Gates, Jr., M.T. Grisaru, M. Roček and W. Siegel: *Superspace*
(Benjamin/Cummings Pub. Co., Reading, Massachusetts, 1983).
- [8] W. Siegel: *Nucl. Phys.* **B238** (1984)307.
- [9] S.J. Gates, Jr. and W. Siegel: *Phys. Lett.* **B206** (1988)631.
- [10] E.M.C. Abreu and C. Wotzasek: *Phys. Rev.* **D58** (1998)101701, hep-th/9805043.
- [11] I.A. Batalin and E.S. Fradkin: *Nucl. Phys.* **B279** (1987)514–528.
- [12] S.J. Gates, Jr. and W. Siegel: *Nucl. Phys.* **B187** (1981)389.
- [13] S.J. Gates, Jr., M.T. Grisaru, M.B. Knutt-Wehlau, M. Roček and O. Solovov:
Phys. Lett. **B396** (1997)167–176;
S.J. Gates, Jr. and S. M. Kuzenko: The CNM-Hypermultiplet Nexus.
hep-th/9810137.
- [14] D.R. Morrison and M.R. Plesser: *Nucl. Phys.* **A46** (Proc. Supl., 1996)177–186,
hep-th/9508107.
- [15] J. Polchinski: TASSI Lectures on D-Branes. hep-th/9611050
- [16] T. Hübsch: *Mod. Phys. Lett.* **A6** (1991)1553–1559.
- [17] W. Siegel: *Physica* **D15**(1985)208–212.
- [18] P.P. Srivastava: *Phys. Rev. Lett.* **63** (1989)2791.
- [19] E. Witten: *J. Diff. Geom.* **17**(1982)661–692.
- [20] E. Witten: *Nucl. Phys.* **B202** (1982)253–316.
- [21] S. Elizur and A. Schwimmer: *Nucl. Phys.* **B226** (1983)109.
- [22] M. Claudson and M.B. Halpern: *Nucl. Phys.* **B250** (1985)689.
- [23] S. Cecotti: *Int. J. Mod. Phys.* **A6** (1991)1749, *Nucl. Phys.* **B355** (1991)755.
- [24] S. Cecotti, L. Girardello and A. Pasquinucci:
Nucl. Phys. **B328** (1989)701, *Int. J. Mod. Phys.* **A6** (1991)2427.
- [25] N. Seiberg and E. Witten: *Nucl. Phys.* **B426** (1994)19–52, Erratum: *ibid.* **B430**
(1994)485–486; *Nucl. Phys.* **B431** (1994)484–550.
- [26] V. Novikov, M. Shifman, A. Vainshtein and V.A. Zakharov: *Nucl. Phys.* **B229**
(1993)381, *Nucl. Phys.* **B229** (1993)407.
- [27] D. Amati, K. Konishi, Y. Meurice, G.C. Rossi and G. Veneziano: *Phys. Rep.* **162C**
(1988)170, and references therein.
- [28] S.J. Gates, Jr. and L. Rana: *Phys. Lett.* **B352** (1995)50–58, hep-th/9504025; *Phys. Lett.* **B369** (1996)262–268, hep-th/9510151.